

**POISON**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY INSTRUCTIONS BEFORE OPENING OR USING**

# OzCrop Imidacloprid 600 FS Insecticide



**ACTIVE CONSTITUENT: 600 g/L IMIDACLOPRID**

|              |           |                    |
|--------------|-----------|--------------------|
| <b>GROUP</b> | <b>4A</b> | <b>INSECTICIDE</b> |
|--------------|-----------|--------------------|



**For the control of various insect pests in a range of crops  
and the protection of spread of barley yellow dwarf virus in  
cereal crops, as specified in the Directions for Use**

**IMPORTANT: READ ATTACHED LEAFLET BEFORE USE**

**CONTENTS 1L, 5L, 10L, 20L, 110L, 200L, 1000L**

**OzCrop Pty Ltd** | Suite 406, 3 Waverley Street, Bondi Junction NSW 2022, Australia  
Tel: (02) 8362 2300

**STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on site. If recycling return clean containers to recycler or point of sale.

If not recycling break, crush, or puncture and bury containers at a local authority landfill. If not available bury the container below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**SAFETY DIRECTIONS**

Harmful if swallowed. Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with the eyes and skin. When opening the container and using the product, wear elbow length PVC gloves. Wash hands after use. After each day's use wash gloves.

**FIRST AID**

If poisoning occurs, contact a Doctor or Poisons Information Centre Phone Australia 13 11 26.

**MATERIAL SAFETY DATA SHEET**

For further information refer to the Material Safety Data Sheet (MSDS), which is available from the supplier.

**MANUFACTURERS WARRANTY AND EXCLUSION OF LIABILITY**

OzCrop Pty Ltd has no control over storage, handling and manner of use of this product. Where this material is not stored, handled or used correctly and in accordance with directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. OzCrop Pty Ltd accepts no responsibility for any loss arising from incorrect storage, handling or use.

APVMA Approval No: 68446 | 58378

|  |   |
|--|---|
| <p>In a Transport Emergency<br/>Dial <b>000</b><br/>Police or Fire Brigade</p> | <p>SPECIALIST ADVICE<br/>IN EMERGENCY ONLY<br/><b>1800 033 111</b><br/>ALL HOURS AUSTRALIA WIDE</p> |
| <p>Toxic 6 diamond</p>   | <p>Pesticide, liquid, toxic N.O.S<br/>(contains imidacloprid)<br/>UN 2902 HAZCHEM 2X</p>            |

Batch No:

Date of Manufacture:

**POISON**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# OzCrop Imidacloprid 600 FS Insecticide

**ACTIVE CONSTITUENT: 250 g/L IMIDACLOPRID**

|              |           |                    |
|--------------|-----------|--------------------|
| <b>GROUP</b> | <b>4A</b> | <b>INSECTICIDE</b> |
|--------------|-----------|--------------------|

**For the control of various insect pests in a range of crops  
and the protection of spread of barley yellow dwarf virus in  
cereal crops, as specified in the Directions for Use**

**THIS LEAFLET IS PART OF THE LABEL**

APVMA Approval No: 68446 | 58378

**OzCrop Pty Ltd** | Suite 406, 3 Waverley Street, Bondi Junction NSW 2022, Australia  
Tel: (02) 8362 2300

**DIRECTIONS FOR USE**

**For ULV application: OzCrop Imidacloprid 600 FS can be bulked up with spraying oils for all uses except those indicated in the critical comments in the Direction for Use table below.**

| Crop  | Pest   | Rate   | Critical Comments  |
|---|--|--|--|
| Cotton  | Thrips   | 875mL or<br>1.17 L/100kg<br>seed   | Thrips damage is dependent upon thrips infesting cotton seedlings and the subsequent growth rate of the plants. Choose a higher rate if thrips pressure is expected (e.g. winter cereal and weeds supporting thrips) and/or cotton seedlings are expected to experience slow growth (e.g. cool weather from early planting or sown in shorter season districts). The mid-rate is considered a general rate for normal conditions.  |
|   | Aphids   | 875mL or<br>1.17 L/100kg<br>seed   | When applied for thrips control, these rates will also control early season aphids.  |
|   | Brown flea beetle  | 875mL or<br>1.17 L/100kg<br>seed   | When applied for thrips control, these rates will also reduce damage to cotyledons caused by brown flea beetle.  |
|   | Wireworms  | 875mL or<br>1.17 L/100kg<br>seed   | Use the higher rate for increased length of control.   |
| Maize,<br>Sorghum,<br>Sunflower,<br>Sweetcorn   | Sugarcane wireworm, ( <i>Agrypnus variabilis</i> ), Eastern false wireworm ( <i>Pterohelaeus darlingensis</i> ) Striate false wireworm ( <i>Pterohelaeus alternatus</i> ), Southern false wireworm ( <i>Gonocephalum macleayi</i> ), Black field earwig ( <i>Nalva lividipes</i> ), Wingless cockroach ( <i>Cosmozosteria</i> spp., <i>Calolampra elegans</i> , <i>C. solidata</i> ) Field cricket ( <i>Teleogryllus commodus</i> ), Black sunflower scarab ( <i>Pseudoheteonyx basicollis</i> ) | Maize:<br>1.4mL/1000<br>seeds<br>Sorghum,<br>sunflower,<br>sweet corn:<br>430mL/100kg<br>of seed | Apply only to high quality seed. Ensure thorough coverage of seed.   |
| Canola  | Aphids<br>Blue Oat mite, red legged<br>earth mite  | 400 mL/<br>100kg of<br>seed  | OzCrop Imidacloprid 600 FS will protect canola seedlings from early season aphid damage.   |
| Forage and<br>seed pasture,<br>e.g. grasses<br>such as<br>ryegrass,<br>fescue and<br>phalaris;<br>clovers such<br>as<br>subterranea<br>n, white, and<br>strawberry;<br>medics;<br>lucerne |  |  | OzCrop Imidacloprid 600 FS will protect emerging seedlings for 3-4 weeks after sowing. Crops must be monitored in the first 4 weeks of emergence to ensure product performance. For autumn sowing: After a pasture phase sow OzCrop Imidacloprid 600 FS treated seed following a well timed spring spray (prior to the development of diapause eggs) of Le-Mat 290SL Insecticide. After a cropping phase a spring spray is not usually required, however if monitoring in spring finds moderate mite populations, a spring spray should be applied. For spring sowing: After a pasture phase apply a spray of omethoate prior to sowing OzCrop Imidacloprid 600 FS treated seed. |
| Forage<br>brassicac   |  |  | OzCrop Imidacloprid 600 FS does not affect the viability of Rhizobia when OzCrop Imidacloprid 600 FS is mixed with inoculant and seed is sown  |

| Crop                             | Pest  | Rate                        | Critical Comments   |
|----------------------------------|---|-----------------------------|---|
| (kale, turnips, rape and Swedes) |   |                             | immediately into moist soil.  |
| Lupins                           |   | 300 mL/100kg seed           |   |
| Cereals                          | Feeding damage caused by wheat aphid and corn aphid | 120 or 240 mL/100kg of seed | Use the higher rate for increased length of control and in areas of high risk.  |
|                                  | Spread of barley yellow dwarf virus                 |                             | In high risk areas with over 500mm of rain or in moderate risk areas with over 400mm of rain in seasons following summer rains, a synthetic pyrethroid top-up spray should be applied 7-8 weeks after sowing. |

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

#### WITHHOLDING PERIOD:

##### Harvest:

**Cotton and food producing crops: NOT REQUIRED WHEN USED AS DIRECTED**

##### Grazing:

**Maize, sorghum, or sweet corn plants growing from treated seed: DO NOT GRAZE OR CUT FOR STOCK FOOD WITHIN 4 WEEKS OF SOWING.**

**Pasture, canola or forage brassica plants grown from treated seed: DO NOT GRAZE OR CUT FOR STOCKFEED FOR SIX WEEKS OF SOWING.**

**Cereal or lupins plants grown from treated seed: DO NOT GRAZE OR CUT FOR STOCK FOOD WITHIN NINE WEEKS OF SOWING.**

#### GENERAL INSTRUCTIONS

#### INSECTICIDE RESISTANCE WARNING

|       |    |             |
|-------|----|-------------|
| GROUP | 4A | INSECTICIDE |
|-------|----|-------------|

For insecticide resistance management OzCrop Imidacloprid 600 FS is a group 4A insecticide. Some naturally occurring insect biotypes resistant to OzCrop Imidacloprid 600 FS and other Group 4A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if OzCrop Imidacloprid 600 FS and other group 4A insecticides are used repeatedly. The effectiveness of OzCrop Imidacloprid 600 FS on resistant individuals could be significantly reduced. Since the occurrence of resistant individuals is difficult to detect prior to use, OzCrop Pty Ltd accepts no liability for any losses that may result from the failure of OzCrop Imidacloprid 600 FS to control resistant insects. OzCrop Imidacloprid 600 FS may be subject to specific resistance management strategies. For further information contact your supplier, OzCrop representative or local agricultural department agronomist.

#### Cotton

##### Mixing and seed treatment

Prior to pouring or pumping OzCrop Imidacloprid 600 FS from the container, either shake the container vigorously or operate pump by-pass to ensure thorough mixing of the contents. Flowable formulations will settle over time. Add most of the water into the mixing vat, followed by flowable fungicides (e.g. quintozone), coating agents (e.g. Peridam, Nancret) and OzCrop Imidacloprid 600 FS and stir until homogeneous. Then add the emulsifiable liquid fungicide (e.g. Apron 200SL) slowly whilst stirring. Calibrate commercial seed dressing equipment to ensure the seed throughput is mixed with the correct dose of slurry seed dressing. Ensure the seed dressing mixture is applied to the maximum number of seeds when initially applied to the seed. Retain seed in mixing chamber until coverage on seed is good. Store treated seed under cover in cool, dry conditions. Do not treat seed with poor viability. Do not store treated seed near foodstuffs or where likely to prove hazardous to humans or animals.

#### Forage and seed pasture, canola, forage brassicas, lupins, summer crops

##### Mixing and seed treatment

Prior to pouring, shake container vigorously, then add the required quantity of OzCrop Imidacloprid 600 FS to sufficient water to give even coverage of the seed to be treated. Place seed in mixing equipment and rotate. Mix OzCrop Imidacloprid 600 FS with sufficient water to give even coverage of seed and spray onto seed. For small seeds, apply a total volume of OzCrop Imidacloprid 600 FS plus water of 1.5 L/100kg seed. For large seeds, apply a total volume of OzCrop Imidacloprid 600 FS plus water of 500 mL/100kg seed. Continue rotation until all liquid is distributed on seed. Store treated seed under cover in cool, dry conditions. Do not treat seed with poor viability. Do not store treated seed near foodstuffs or where likely to prove hazardous to humans or animals. Do not carry over maize, sorghum, sunflower or sweet corn seed from one season to the next season. The insecticidal activity of OzCrop Imidacloprid 600 FS on the seed is maintained for a minimum of two seasons. Flow rate: OzCrop Imidacloprid 600 FS treated seed may slow flow rate of seed. Check flow rate of seed through sowing machinery before sowing to ensure the desired seeding rate is achieved.

### **Cereal crops**

#### **Mixing and seed treatment**

Prior to pouring, shake container vigorously, then add the required quantity of OzCrop Imidacloprid 600 FS to sufficient water to give even coverage of the cereal seed to be treated. The quantity of water used for mixing will vary depending on type of equipment and quantity of seed. Use a minimum of 400 mL of mixture (product + water) with each 100 kg of seed. Whatever dilution is used it is essential that 120 mL or (240 mL) of OzCrop Imidacloprid 600 FS is applied/100 kg seed. The mixture should be gently stirred regularly.

#### **Seed quality**

OzCrop Imidacloprid 600 FS seed treatment should not be used on seed with more than 12% moisture content, or on sprung, sprouted, damaged or severely pinched seed of poor viability. If in doubt, have a germination test carried out on the seed before treatment to ensure that it is of acceptable standard. Treating with OzCrop Imidacloprid 600 FS as directed may raise the moisture level of the seed by up to 0.6%, depending on conditions at treatment. The use of OzCrop Imidacloprid 600 FS mixed with the water at recommended rates will have no effect on the storage life of treated sound seed.

#### **Compatibility**

OzCrop Imidacloprid 600 FS is compatible with Baytan C Flowable Seed Dressing and Raxil C Flowable Seed Dressing for use on cereal seeds. Do not use more than 600 mL of total mixture /100 kg seed (i.e. OzCrop Imidacloprid 600 FS plus Baytan or Raxil plus water). OzCrop Imidacloprid 600 FS may be applied with Jockey Systemic Seed Fungicide or with Semevin P Seed Dressing Insecticide.

#### **Storage of treated seed**

Do not store treated seed near foodstuffs or where likely to prove hazardous to humans or animals. If the seed is not used immediately after treatment it should be stored in a dry, well ventilated place. Although OzCrop Imidacloprid 600 FS has no effect on the viability of treated seed, subsequent germination can be adversely affected by poor storage conditions such as high moisture combined with high temperatures. No liability can therefore be accepted for the performance of stored treated seed.

### **PRECAUTIONS**

When treated seed is stored it should be kept apart from other grain and the bags or other containers should be clearly marked to indicate the contents have been treated with this product. Do NOT allow seed treated with this product to contaminate seed intended for human consumption. Do NOT use treated seed for human consumption. Bags which have held treated seed are not to be used for any other purpose.

### **PROTECTION OF LIVESTOCK**

Seed treated with this product must not be used for animal consumption or poultry feed or mixed with animal feed. Do NOT allow seed treated with this product to contaminate seed intended for animal consumption.

### **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

Do NOT contaminate ponds, waterways and drains with this product, used containers or bags which have held treated seed. Do NOT feed treated seed or otherwise expose to wild or domestic birds. Any

spillages of treated seed, however minor, must be cleaned up immediately, preferable by recovery and re-use. If disposal is required, ensure treated seeds as thoroughly buried and not accessible to birds and other wildlife.

#### **STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on site. If recycling return clean containers to recycler or point of sale.

If not recycling break, crush, or puncture and bury containers at a local authority landfill. If not available bury the container below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

#### **SAFETY DIRECTIONS**

Harmful if swallowed. Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with the eyes and skin. When opening the container and using the product, wear elbow length chemical resistant gloves. Wash hands after use. After each day's use wash gloves.

#### **MATERIAL SAFETY DATA SHEET**

For further information refer to the Material Safety Data Sheet (MSDS), which is available from the supplier.

#### **MANUFACTURERS WARRANTY AND EXCLUSION OF LIABILITY**

OzCrop Pty Ltd has no control over storage, handling and manner of use of this product. Where this material is not stored, handled or used correctly and in accordance with directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. OzCrop Pty Ltd accepts no responsibility for any loss arising from incorrect storage, handling or use.

|  |   |
|--|---|
| <p>In a Transport Emergency<br/>Dial <b>000</b><br/>Police or Fire Brigade</p> | <p>SPECIALIST ADVICE<br/>IN EMERGENCY ONLY<br/><b>1800 033 111</b><br/>ALL HOURS AUSTRALIA WIDE</p> |
| <p>Toxic 6 diamond</p>   | <p>Pesticide, liquid, toxic N.O.S<br/>(contains imidacloprid)<br/>UN 2902 HAZCHEM 2X</p>            |