

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

 OzCrop

Azoxycypro WG

FUNGICIDE

ACTIVE CONSTITUENTS: 600G/KG AZOXYSTROBIN
240G/KG CYPROCONAZOLE

GROUP **3 | 11** FUNGICIDE

*Controls certain fungal diseases in Barley and Wheat
as specified in the Directions for Use*

NET CONTENTS:
500g-20Kg

NOT A DANGEROUS GOOD
ACCORDING TO THE AUSTRALIAN
DANGEROUS GOODS (ADG) CODE.

BATCH NO.
DATE OF MANUFACTURE:

OzCrop Pty Ltd ABN 16 160 656 431
Suite G13/25 Solent Circuit, Norwest, NSW 2153
Tel: 02 8123 0170 • Fax: 02 8123 0171
Email: enquiries@ozcrop.com.au
www.ozcrop.com.au

DIRECTIONS FOR USE RESTRAINTS

DO NOT apply by aircraft.

DO NOT apply more than 2 applications of OZCROP AZOXYCYPRO WG FUNGICIDE (or any other strobilurin or Group 3 fungicide) in any one season on the same paddock.

DO NOT exceed a combined total of 267 g/ha of OZCROP AZOXYCYPRO WG FUNGICIDE in any one season on the same paddock.

SPRAY DRIFT RESTRAINTS

DO NOT apply when wind speed is less than 3 or more than 20 kilometres per hour, as measured at the application site.

DO NOT apply during surface temperature inversion conditions at the application site.

Users of this product MUST make an accurate written record of the details of each spray application within 24 hours following application, and must KEEP this record for at least 2 years. The spray application details that must be recorded are: 1 date with start and finish times of application 2 location address and paddock(s) sprayed 3 full name of this product 4 amount of product used per hectare and number of hectares applied to 5 crop or situation and weed or pest 6 wind speed and direction during application 7 air temperature and relative humidity during application 8 nozzle brand, type, spray angle, nozzle capacity and spray system pressure measured during application 9 name and address of person applying this product. (Additional record details may be required by the state or territory where this product is used.)

MANDATORY NO-SPRAY ZONES

DO NOT apply if there are aquatic and wetland areas including aquaculture ponds, surface streams and rivers within 50 metres downwind from the application area.

Crop	Diseases	Rate/ha	Critical Comments
Barley	Leaf Rust (Puccinia hordei) Net blotch: Net Form (Pyrenophora teres f. teres) Spot Form (Pyrenophora teres f. maculata)	67-267 g	Apply from jointing (Z30) and before infection averages 10% on the flag-2 leaf. Ensure thorough coverage of stems and leaves. Use higher rates under high infection pressure or when longer residual protection is required. Lower rates are effective under low disease pressure but have reduced residual effect. Repeat application at early booting stage if necessary. Two applications of the lower rate will generally provide better control than a single application at the higher rate. Addition of ADIGOR® at 2 % v/v improves disease control, especially at the lower rate.
	Powdery Mildew (Blumeria graminis hordei)	133-267 g	Spray at the first sign of the disease during the tillering stage. A repeat spray 21 to 28 days later may be required. Ensure thorough coverage of stems and leaves. Higher rates provide longer protection.
Wheat	Leaf Rust (Puccinia triticina)	133-267 g	Apply after flag leaf is 70% emerged and before disease levels reach 1% of flag leaf area. Consider control if disease is greater than 5 to 10% on any lower leaf layer. Use higher rates under high infection pressure or when longer residual protection is required. Lower rates are effective under low disease pressure but have reduced residual effect. Ensure thorough coverage.
	Powdery Mildew (Blumeria graminis tritici)		Spray at the first sign of the disease during the tillering stage. A repeat spray 21 to 28 days later may be required. Ensure thorough coverage of stems and leaves. Higher rates provide longer protection.
	Stripe Rust (Puccinia striiformis)		Spray at the first sign of the disease and before 10 to 20% of leaves are infected. A repeat spray 21 to 28 days later may be required. Use higher rates under high infection pressure or when longer residual protection is required.
	Yellow Spot (Pyrenophora tritici-repentis)		Apply once between 70% flag leaf emergence and early flowering. Use higher rates under high infection pressure or where longer residual protection is required. Apply after 70% flag leaf emergence and before disease levels reach 5% on flag leaf. Higher rates give longer residual protection and often better economic returns.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS

Barley Wheat:

Grazing: DO NOT GRAZE OR CUT FOR STOCK FEED FOR 21 DAYS AFTER APPLICATION

Harvest: DO NOT HARVEST FOR 6 WEEKS AFTER APPLICATION

EXPORT SLAUGHTER INTERVAL

DO NOT slaughter animals destined for export within 7 days of consumption of treated cereal forage or straw.

WARNING

Azoxytrobin is extremely phytotoxic to certain apple varieties.

AVOID SPRAY DRIFT. Extreme care must be used to prevent injury to apple trees. DO NOT spray OZCROP AZOXYCYPRO WG FUNGICIDE where spray drift may reach apple trees.

DO NOT spray when conditions favour drift beyond the area intended for application. Conditions which may contribute to drift include thermal inversions, excessive wind speed, certain spray nozzle/pressure combinations, small spray droplet size, etc.

DO NOT use spray equipment which has been previously used to apply OZCROP AZOXYCYPRO WG FUNGICIDE to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity.

GENERAL INSTRUCTIONS

MIXING

OZCROP AZOXYCYPRO WG FUNGICIDE is a water dispersible granule formulation that mixes readily with water and is applied as a spray.

1. Partly fill the spray tank with water
2. Start the agitation

3. Add the correct amount of product to the spray tank with the agitation system running
4. Continue agitation while topping up the tank with water and while spraying
5. Use the spray mix within 24 hours of preparation

GROWING CONDITIONS

In general OZCROP AZOXYCYPRO WG FUNGICIDE should be applied under good growing conditions with adequate soil moisture. OZCROP AZOXYCYPRO WG FUNGICIDE should not be applied to crops that are under stress due to very dry, very wet, frosty conditions, nutrient deficiency or high insect pressure, as this may give less than reliable results.

Spray Timing

Always inspect crops to assess disease development immediately before spraying. Best results will be achieved from applications made in the earliest stages of disease development or as a protectant treatment following a disease risk assessment.

Applications of OZCROP AZOXYCYPRO WG FUNGICIDE should be made within the crop growth stages ranging from tillering to early flowering (Z26 to Z62). The object of spraying is to keep the upper 2 to 3 leaves green and functioning through grain filling stage.

APPLICATION

Apply by boom spray and ensure complete coverage of all leaves and stem is obtained. As a guide, apply in spray volumes ranging from 50 to 100 L water per hectare. The use of flat fan nozzles is recommended using a fine to medium quality spray pattern, with pressures in the range 100 to 400 kPa.

COMPATIBILITY/TANK MIXING

OZCROP AZOXYCYPRO WG FUNGICIDE is physically compatible with a wide range of products, however the biological compatibility of these mixtures may not have been fully tested under all environmental and biological conditions. A mixture of OZCROP AZOXYCYPRO WG FUNGICIDE with any other product may in some circumstances be ineffective or may cause damage. If tank mixes are to be used observe all directions, precautions and limitations on all products to be used. As formulations of other manufacturer's products are beyond the control of Rainbow, and the quality of water may vary with location, all mixtures should be tested prior to mixing commercial quantities.

FUNGICIDE RESISTANCE WARNING

OZCROP AZOXYCYPRO WG FUNGICIDE is a combination of a DMI and a strobilurin and related products fungicide. For fungicide resistance management OZCROP AZOXYCYPRO WG FUNGICIDE is both a Group 3 and a Group 11 fungicide. Some naturally occurring individual fungi resistant to OZCROP AZOXYCYPRO WG FUNGICIDE and other Group 3 and/or Group 11 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by OZCROP AZOXYCYPRO WG FUNGICIDE or other Group 3 and/or Group 11 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, OzCrop Pty Ltd accepts no liability for any losses that may result from the failure of OZCROP AZOXYCYPRO WG FUNGICIDE to control resistant fungi.

GROUP 3 | 11 FUNGICIDE

PRECAUTIONS

Re-entry Period: DO NOT allow entry into treated areas until the spray has dried unless wearing cotton overalls (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF WILDLIFE, FISH CRUSTACEANS AND ENVIRONMENT

Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.

STORAGE AND DISPOSAL

Store in closed original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

Poisonous if swallowed. May irritate eyes. Avoid contact with eyes. When opening the container and preparing spray and if applying by boomspray equipment, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and rubber gloves. Wash hands after use. After each day's use wash gloves and contaminated clothing.

Safe Work Australia:

Harmful if inhaled.

Very toxic to aquatic life with long lasting effects.

Use only outdoors or in a well-ventilated area.

Avoid breathing dust/fumes.

Do not eat, drink or smoke when using this product.

Avoid release to the environment.

IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing.

Rinse mouth.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. If in eyes wash out immediately with water.

SAFETY DATA SHEET

If additional hazard information is required refer to the Safety Data Sheet. For a copy phone 07 3198 4930.

DISCLAIMER

This product complies with the specifications in its statutory registration. Implied terms and warranties are excluded. OzCrop liability for breach of the express or any non-excludable implied warranty is limited to product replacement or purchase price refund. The purchaser must determine suitability for intended purpose and take all proper precautions in the handling, storage and use of the product including those on the label and/or safety data sheet failing which OzCrop Pty Ltd shall have no liability.



OzCrop Pty Ltd ABN 16 160 656 431

Suite G13/25 Solent Circuit, Norwest, NSW 2153

Tel: 02 8123 0170 • Fax: 02 8123 0171

Email: enquiries@ozcrop.com.au

visit our website www.ozcrop.com.au

APVMA Approval No.: 87237/117848

Label Release: 201906

