

MAIN PANEL

Text above this line is not part of the label

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING


OzCrop
BROMOXYNIL 200
SELECTIVE HERBICIDE

ACTIVE CONSTITUENT: 200g/L BROMOXYNIL present as the n-octanoyl ester
SOLVENT: 610g/L LIQUID HYDROCARBON

GROUP	C	HERBICIDE
-------	----------	-----------

For the control of certain broadleaf weeds in cereals, linseed, lucerne, pastures and turf as specified in the Directions for Use table.

**IMPORTANT: READ THE ATTACHED LEAFLET
BEFORE USING THIS PRODUCT**



Contents: 5L, 20L, 110L, 200L & 1000L

IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE
EMERGENCY CONTACT 1800 033 111 AUSTRALIA WIDE, 24 HOURS

OzCrop Pty Ltd | Suite 406, 3 Waverley Street, Bondi Junction NSW 2022, Australia
Tel: (02) 8362 2300

APVMA Approval No. 68441 / 58372

MAIN PANEL

Text above this line is not part of the label

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Protect from frost. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury the empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

Container - Micro Matic Valve (110L)

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Protect from frost. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the container with water or any other foreign matter. After each use of the product please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the container have been used, please return the empty container to the point of purchase.

Refillable containers (1000L only)

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Protect from frost. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Product is poisonous if inhaled or swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing spray wear elbow-length PVC gloves and face shield. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26). If swallowed, DO NOT induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS) which can be obtained from the supplier.

NOTICE TO BUYER

OzCrop Pty LTD has no control over storage, handling and manner of use of this product. Where this material is not stored, handled or used correctly and in accordance with directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. OzCrop Pty LTD accepts no responsibility for any loss arising from incorrect storage, handling or use.

Batch No.:

Date of Manufacture

LABEL LEAFLET

Text above this line is not part of the label

POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

OzCrop BROMOXYNIL 200 SELECTIVE HERBICIDE

ACTIVE CONSTITUENT: 200g/L BROMOXYNIL present as the n-octanoyl ester

SOLVENT: 610g/L LIQUID HYDROCARBON

GROUP	C	HERBICIDE
-------	----------	-----------

For the control of certain broadleaf weeds in cereals, linseed, lucerne, pastures and turf as specified in the Directions for Use table.

**IMPORTANT: READ THIS LEAFLET
BEFORE USING THIS PRODUCT**

Contents: 5L, 20L, 110L, 200L & 1000L

IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE
EMERGENCY CONTACT 1800 033 111 AUSTRALIA WIDE, 24 HOURS

OzCrop Pty Ltd | Suite 406, 3 Waverley Street, Bondi Junction NSW 2022, Australia
Tel: (02) 8362 2300

APVMA Approval No. 68441 / 58372

LABEL LEAFLET

Text above this line is not part of the label

DIRECTIONS FOR USE:

RESTRAINTS:

DO NOT spray if crops or weeds are stressed due to dry or excessively moist conditions.

DO NOT apply to diseased or frost-affected crops or if frosts are imminent.

DO NOT apply if rain is expected within three hours.

SITUATION & CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE/HA	CRITICAL COMMENTS
Wheat, Barley, Cereal Rye, Oats and Triticale: including those undersown with Clover, Lucerne or Medic (Vic only);	Amsinckia (<i>Amsinckia</i> spp.), Black Bindweed (Climbing Buckwheat) (<i>Polygonum convolvulus</i>), Capeweed (<i>Arctotheca Calendula</i>), Chamomile (<i>Matricaria matricarioides</i>), Common Peppergrass (<i>Lepidium hyssopifolium</i>), Corn Gromwell (Ironweed, Sheepweed) (<i>Buglossoides arvensis</i>), Fat Hen (<i>Chenopodium album</i>), Field Madder (<i>Sherardia arvensis</i>), Lesser Swinecress (<i>Coronopus didymus</i>), Mountain Sorrel (<i>Calandrinia</i> spp.), Saffron Thistle (<i>Carthamus Lanatus</i>), Shepherd's Purse (<i>Capsella bursa-pastoris</i>), Tree Hogweed (<i>Polygonum patulum</i>), Variegated Thistle (<i>Silybum marianum</i>)	Up to the 4 leaf stage or when plants are no more than 35 mm in diameter	All States	1.4 L	CROP STAGE: Wheat, Barley, Cereal, Rye, Oats and Triticale: 3 leaf to fully tillered (Zadoks Scale Z13-30). Linseed: 50 to 150 mm high. Lucerne and Clover: After seedlings have one trifoliate leaf or before there is 150 mm regrowth in established stands. Lotus: Apply post-emergence at least 4 weeks after sowing (crop seedlings with a min. of 2-6 true leaves). APPLICATION: Avoid application during temperatures above 20°C as seedling mortality may occur and established plants may be damaged. Good coverage of the spray on the weeds is essential. Apply only in the Autumn and Winter (mid-April to end of August). For optimum control, apply only when the weeds are actively growing and before the weeds are shielded by the crop. CAUTION: Lucerne and Clover: This product will cause slight leaf burning of undersown lucerne and clover. DO NOT apply to Persian clover or Berseem clover. Linseed: DO NOT spray linseed by aerial application. Lotus: Some initial scorching of leaves may occur in some cases. Medics: Application to some cultivars of Medics may result in unacceptable crop damage in some situations. ^{#1} Where Wireweed is growing under stress, it has been found to be less susceptible. ^{#2} Effective controls of Patersons Curse and Fumitories will only be obtained under good growing conditions.
Linseed; Clover or Lucerne Pastures; Lotus seed or forage crops (Qld only)	Mexican Poppy (<i>Argemone mexicana</i>), Threecornered Jack (<i>Emex australis</i>) (Doublegee, Spiny Emex), Turnip Weed (<i>Rapistrum rugosum</i>) (Giant Mustard), Wild Radish (<i>Raphanus raphanistrum</i>), Wild Mustard (<i>Sisymbrium</i> spp.), Wild Turnip (<i>Brassica tournefortii</i>), Wireweed ^{#1} (<i>Polygonum aviculare</i>), Patersons Curse ^{#2} (<i>Echium plantagineum</i>) (Salvation Jane)	Up to the 6 leaf stage or when plants are no more than 50 mm in diameter		2 L	
	Fumitories ^{#2} (<i>Fumaria</i> spp.)	Up to the 4 leaf stage or when plants are no more than 50 mm in diameter	Qld, NSW, Vic, SA,		

LABEL LEAFLET

Text above this line is not part of the label

Wheat, Barley, Cereal Rye, Oats and Triticale: including those undersown with Clover, Lucerne or Medic (Vic only); Linseed; Clover or Lucerne Pastures; Lotus seed or forage crops (Qld only)	Adonis (<i>Adonis microcarpa</i>) (Pheasant's Eye), Birds Eye (<i>Cotula australis</i>) (Carrot Weed)	Up to the 4 true leaf stage	WA only SA only	700mL	DO NOT use where undersown medics occur.
		Greater than the 4 true leaf stage.		1.4 L	
	Fireweed (<i>Senecio</i> spp.)	Prior to flower development	Qld, NSW only		Apply during the Autumn-Winter period when weed are young and actively growing.
		Early flowering stage		2.8 L	
Wheat, Barley, Cereal Rye, Oats and Triticale: including those undersown with Clover, Lucerne or Medic Clover Seed Crops	Amsinckia (<i>Amsinckia</i> spp.), Capeweed (<i>Arctotheca calendula</i>), Chamomile (<i>Matricaria matricarioides</i>), Charlock (<i>Sinapis arvensis</i>), Corn Gromwell (<i>Buglossoides arvensis</i>) (Ironweed, Sheepweed), Fat Hen (<i>Chenopodium album</i>), Field Madder (<i>Sherardia arvensis</i>), Hexham Scent (<i>Melilotus indicus</i>) (King Island Melilot), Horned Poppy (<i>Glaucium flavum</i>), Indian Hedge Mustard (<i>Sisymbrium orientale</i>), Lesser Swinecress (<i>Coronopus didymus</i>), Mexican Poppy (<i>Argemone mexicana</i>), Mintweed (<i>Salvia reflexa</i>), Mountain Sorrel (<i>Calandrinia</i> spp.) (Purple Calandrinia), Patersons Curse (<i>Echium plantagineum</i>) (Salvation Jane), Rough Poppy (<i>Papaver hybridum</i>), Saffron Thistle (<i>Carthamus lanatus</i>), Shepherd's Purse (<i>Capsella bursa-pastoris</i>), Slender Thistle (<i>Carduus tenuiflorus</i>), Turnip Weed (<i>Rapistrum rugosum</i>) (Giant Mustard), Wild Radish (<i>Raphanus raphanistrum</i>)	Up to the 5 leaf stage or when weed plants are no more than 50mm in diameter	WA only	750mL PLUS 550mL of 500g/L 2,4-D amine OR PLUS 700mL of 500g/L MCPA amine	CROP STAGE: CEREALS: 3 leaf to fully tillered. (Zadoks Scale Z13-30). CLOVER: Apply after clover seedlings have at least three trifoliate leaves. APPLICATION: Avoid application during temperatures above 20°C as seedling mortality may occur and established plants may be damaged. Apply in Autumn and Winter (mid-April to end of August). Apply only when weeds are actively growing and before the weeds are shielded by the crop. CAUTION: Some crop damage and/or yield loss may occur in Olympic and Shortim wheat cultivars. DO NOT apply to Persian clover or Berseem clover. A slight burning of clover may occur. DO NOT use 2,4-D or MCPA tank mix if wheat, oats, barley or triticale are undersown with lucerne, clover or medics unless some damage is acceptable. This product may be tank mixed with Diclofop 375 Selective Herbicide for control of Annual Ryegrass and Wild Oats in Wheat and Barley only. DO NOT use 2,4-D or MCPA tank mix if using a tank mix with Diclofop.
			SA only	1.4 L PLUS 550mL of 500g/L 2,4-D amine OR PLUS 700mL of 500g/L MCPA amine	

LABEL LEAFLET

Text above this line is not part of the label

SITUATION & CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE/ HA	CRITICAL COMMENTS
Wheat, Barley, Cereal Rye, Oats and Triticale: including those undersown with Clover, Lucerne or Medic Clover Seed Crops (continued)	Common Peppergrass (<i>Lepidium hyssopifolium</i>), Tree Hogweed (<i>Polygonum patulum</i>)	Up to the 5 leaf stage or when weed plants are no more than 50 mm in diameter	Vic only	1.4 L PLUS 300mL of 500g/L MCPA amine	CROP STAGE: CEREALS: 3 leaf to fully tillered. (Zadoks Scale Z13-30). CLOVER: Apply after clover seedlings have at least three trifoliate leaves. APPLICATION: Avoid application during temperatures above 20°C as seedling mortality may occur and established plants may be damaged. Apply in Autumn and Winter (mid-April to end of August). Apply only when weeds are actively growing and before the weeds are shielded by the crop. CAUTION: Some crop damage and/or yield loss may occur in Olympic and Shortim wheat cultivars. DO NOT apply to Persian clover or Berseem clover. A slight burning of clover may occur. DO NOT use 2,4-D or MCPA tank mix if wheat, oats, barley or triticale are undersown with lucerne, clover or medics unless some damage is acceptable. This product may be tank mixed with Diklofop for control of Annual Ryegrass and Wild Oats in Wheat and Barley only. DO NOT use 2,4-D or MCPA tank mix if using a tank mix with Diklofop.
	Black Bindweed (<i>Polygonum convolvulus</i>) (Climbing Buckwheat)	Up to the 8 leaf stage	Vic, SA only		
Wheat, Barley, Oats and Triticale	Spiny Emex (<i>Emex australis</i>) (Doublegee, Threecornered Jack), Wireweed (<i>Polygonum aviculare</i>) Variegated Thistle (<i>Silybum marianum</i>) Fumitories (<i>Fumaria</i> spp.)	Up to the 4 leaf stage or when weed plants are no more than 50 mm in diameter	Qld, NSW, Vic, Tas, SA only	1.4 L PLUS 300mL of 500g/L MCPA amine	CROP STAGE: 5 leaf to fully tillered (Zadoks scale Z15-30); Apply under good growing conditions. BOOM SPRAYING: Use a minimum of 220 L spray per hectare on Black Bindweed when crop is greater than 300 mm high. CAUTION: Some crop damage and/or yield loss may occur in Olympic and Shortim wheat cultivars. DO NOT mix with 2,4-D or MCPA tank mix if wheat, oats, barley or triticale are undersown with lucerne, clover or medics unless some damage is acceptable. DO NOT apply to medics (Western Australia only). This product may
	Wild Mustard (<i>Sisymbrium</i> spp.), Wild Turnip (<i>Brassica tournefortii</i>)		Qld, NSW, Vic only		
	Spiny Emex (<i>Emex australis</i>) (Doublegee, Threecornered Jack), Wireweed (<i>Polygonum aviculare</i>)		WA only		
	Fumitories (<i>Fumaria</i> spp.)	4 to 5 leaf stage	Qld, NSW, Vic, SA only	2.1 L plus 550mL of	

LABEL LEAFLET

Text above this line is not part of the label

	Black Bindweed (<i>Polygonum convolvulus</i>) (Climbing Buckwheat), Chamomile (<i>Matricaria matricarioides</i>), Fat Hen (<i>Chemopodium album</i>), Field Madder (<i>Sherardia arvensis</i>), Lesser Swinecress (<i>Coronopus didymus</i>), Mountain Sorrel (<i>Calandrinia menziesii</i>), Patersons Curse (<i>Echium plant agineum</i>) (Salvation Jane), Shepherd's Purse (<i>Capsella bursa-pastoris</i>), Variegated Thistle (<i>Silybum marianum</i>)	5 to 8 leaf stage	All States	500g/L 2,4-D amine or plus 700mL of 500g/L MCPA amine	be tank mixed with Diclofop for control of Annual Ryegrass and Wild Oats in Wheat and Barley only. DO NOT use 2,4-D or MCPA tank mix if using a tank with Diclofop.
Pastures, Roadsides and Rights- of- Way	African Daisy (<i>Senecio pterophorus</i>)	Up to and including the flowering stage	SA only	300mL per 100 L water	The plants, including the stems, must be thoroughly wet to the point of run-off. Apply when the plants are actively growing. A non- ionic wetter should be added at 175 mL per 100 litres of spray. A follow-up treatment may be necessary for large, well- established plants.
	Boneseed (<i>Chrysanthemoides monilifera</i>)	Seedling only	Vic, Tas only	160mL per 100 L water	Ensure an overall spray coverage on weed seedlings.

SITUATION & CROP	WEEDS CONTROLLED	STAT E	RATE			CRITICAL COMMENTS
Turf: Established Couch (including Queensland Blue), Buffalo, Bent, Paspalum, Kikuyu rescue and Rye Lawns	Bindy-eye (<i>Calotis hispidula</i>), Jo-jo (<i>Onehunga</i>) (<i>Soliva pterospema</i>), Cudweed (<i>Gnaphalium</i> spp.)	All States	6.0L/ 500L water per ha	High Volume: 300mL per 100L	60mL/ 100m ² per 15L knapsack	Apply evenly over the area to be treated. A slight browning of grasses, especially Paspalum, may occur in hot conditions.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS
LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

WITHHOLDING PERIODS:

**CEREALS, PASTURES AND FORAGE CROPS: DO NOT GRAZE OR CUT FOR STOCK
FOOD FOR 8 WEEKS AFTER APPLICATION.**

GENERAL INSTRUCTIONS

RESISTANT WEEDS WARNING

GROUP	C	HERBICIDE
-------	----------	-----------

OzCrop Bromoxynil 200 Selective Herbicide is a member of the Nitrite group of herbicides. OzCrop Bromoxynil 200 has the photosystem II inhibitor mode of action. For weed resistance management OzCrop Bromoxynil 200 is a Group C herbicide.

LABEL LEAFLET

Text above this line is not part of the label

Some naturally-occurring weed biotypes resistant to OzCrop Bromoxynil 200 and other Group C herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by OzCrop Bromoxynil 200 or other Group C herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, OzCrop Pty Ltd. accepts no liability for any losses that may result from the failure of OzCrop Bromoxynil 200 to control resistant weeds. However resistance should not be assumed without first viewing the method of application, the coverage and the timing of application.

MIXING

To ensure even mixing, half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly, then add the remainder of the water. Agitate again before spraying commences.

APPLICATION

Boom Sprayer: Apply 50 to 200 (WA: 50 to 100) litres of water per hectare. Calibrate the machine before use to ensure that the correct application rate is obtained. Check height of boom above the crop to ensure that the spray is evenly distributed. A minimum of 160 litres and preferably 200 litres of water per hectare is recommended where the weed infestation is heavy or the crop cover is dense. Wash the sprayer thoroughly after use with clean water.

Aerial Application: Apply a minimum of 22 litres of water per hectare. Markers should always move upwind before the aircraft passes overhead. Application by microneaire equipment may give variable results.

COMPATIBILITY

This product may be mixed in the spray tank with one of the following herbicides or insecticides according to the directions of use of this product: Chlorpyrifos, Dicer 500 (Dicamba 500), Dimethoate, Diklofop 375 (Diclofop-Methyl 375), MCPA, Omethoate and Putra-amine 625 (2,4-D Amine 625).

As formulations of other manufactures' products are beyond the control of OzCrop Pty Ltd., all mixtures should be tested prior to mixing commercial quantities.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

Dangerous to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Protect from frost. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury the empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

LABEL LEAFLET

Text above this line is not part of the label

Container - Micro Matic Valve (110L)

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Protect from frost. **DO NOT** tamper with the Micro Matic valve or the security seal. **DO NOT** contaminate the container with water or any other foreign matter. After each use of the product please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the container have been used, please return the empty Envirodrum to the point of purchase.

Refillable containers (1000L only)

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Protect from frost. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Product is poisonous if inhaled or swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing spray wear elbow-length PVC gloves and face shield. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26). If swallowed, DO NOT induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS) which can be obtained from the supplier.

NOTICE TO BUYER

OzCrop Pty LTD has no control over storage, handling and manner of use of this product. Where this material is not stored, handled or used correctly and in accordance with directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. OzCrop Pty LTD accepts no responsibility for any loss arising from incorrect storage, handling or use.

IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE
EMERGENCY CONTACT 1800 033 111 AUSTRALIA WIDE, 24 HOURS