

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

OzCrop

Cyprofludox WG

FUNGICIDE

ACTIVE CONSTITUENTS: 375 g/kg CYPRODINIL
250 g/kg FLUDIOXONIL

GROUP **9|12** FUNGICIDE

For the control of various fungal diseases in Capsicum, Cut Flowers, Grapes, Leafy Vegetables, Lettuce, Nursery Stock, Onions, Pyrethrum and Strawberries.

NET CONTENTS:

1 kg - 20 kg

NOT A DANGEROUS GOOD
ACCORDING TO THE AUSTRALIAN
DANGEROUS GOODS (ADG) CODE.

BATCH NO.

DATE OF MANUFACTURE:

OzCrop Pty Ltd ABN 16 160 656 431
G13/25 Solent Circuit, NORWEST NSW 2153
Telephone: (02) 8123 0170 • Facsimile: (02) 8123 0171
Email: enquiries@ozcrop.com.au
www.ozcrop.com.au

DIRECTIONS FOR USE

Restraints

DO NOT apply by aircraft

DO NOT apply by equipment carried on the back of the user.

Spray Drift Restraints

Except when applying with orchard airblast equipment, DO NOT apply with spray droplets smaller than a MEDIUM spray droplet size category according to nozzle manufacturer specifications that refer to ASAE S572 Standard or the BCPC Guideline.

DO NOT apply when wind speed is less than 3 or more than 20 kilometres per hour as measured at the application site.

DO NOT apply during surface temperature inversion conditions at the application site.

Users of this product MUST make an accurate written record of the details of each spray application within 24 hours following application and KEEP this record for a minimum of 2 years. The spray application details that must be recorded are:

- date with start and finish times of application;
- location address and paddock/s sprayed;
- full name of this product;
- amount of product used per hectare and number of hectares applied to; crop/situation and weed/pest;
- wind speed and direction during application;
- air temperature and humidity;
- nozzle brand, type, spray angle, nozzle capacity and spray system pressure measured during application;
- name and address of person applying this product. (Additional record details may be required by the state or territory where this product is used).

Mandatory No-Spray Zones

DO NOT apply if there are aquatic and wetland areas including aquacultural ponds, surface streams and rivers within 30 metres downwind from the application area.

CROP	PEST	RATE/Ha	CRITICAL COMMENTS
Capsicum	Botrytis rot (<i>Botrytis cinerea</i>), Sclerotinia rot (<i>S. minor</i> or <i>S. sclerotiorum</i>)	100 g/100 L or 1 kg/ha	Apply prior to or at onset of disease development. DO NOT apply more than 2 applications per crop per season with an interval of 7 to 14 days between applications. Use the shorter spray interval under high disease pressure or when continual wet, cloudy conditions favour disease development.
Cut Flowers	Botrytis rot	80 g/100 L or 800 g/ha plus 0.02% non-ionic surfactant	Monitor environmental conditions to determine time of application. Apply before disease appears when conditions favour infection. Apply by low or high volume spray. Apply a maximum of 2 applications within an annual production cycle. For resistance management this product should form part of a disease management strategy. Crop Damage This product has demonstrated toxicity under certain situations, application methods, applications rates and weather conditions to nursery stocks including Impatiens, New Guinea Impatiens, Geraniums, Ferns, Anemone, Begonia, Kalanchoe, Saint Paulia and Exacum affine. This product may also leave undesirable spray residue on foliage and flowers. It is recommended that users treat a sample number of plants or a small area prior to whole crop treatment and monitor any phytotoxic effects that will compromise production goals. This action cannot guarantee crop safety as application, environmental and crop conditions may vary from test treatment to whole crop treatment.
Grapes (except grapes grown for dried fruit production)	Grey Mould (Bunch Rot) <i>Botrytis cinerea</i>	Dilute Spraying 80 g/100 L Concentrate Spraying 800 g/ha	DO NOT apply more than 2 applications of OzCrop Cyprofludox WG Fungicide per season. Applications can be made at 5% capfall, 80% capfall, pre-bunch closure or veraison. Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. DO NOT use at rates greater than 200 g/100 L of water. This use is subject to a CropLife fungicide resistance management strategy (see Resistance Management).

CROP	PEST	RATE/Ha	CRITICAL COMMENTS
Leafy vegetables (including spinach, silver beet, kale, endive, mustard, cress, chard, rocket, Asian leafy greens and Chinese cabbage)	<i>Sclerotinia (Sclerotinia minor and Sclerotinia sclerotiorum)</i>	800g to 1 kg/ha	Apply OzCrop Cyprofludox WG Fungicide prior to or at the onset of disease. Apply a second application 7 to 14 days later if conditions continue to remain favourable for disease development. Use a 7 day spray interval and high rate under high disease pressure or where conditions are conducive for disease development. DO NOT apply more than 2 applications of OzCrop Cyprofludox WG Fungicide per crop.
Lettuce	<i>Sclerotinia Rot (Lettuce Drop) Sclerotinia minor and Sclerotinia sclerotiorum</i>		Apply OzCrop Cyprofludox WG Fungicide prior to or at the onset of disease. Best results are achieved using a preventative control program, with the first spray commencing soon after transplanting on a 7 to 14 spray interval. Use a 7 day spray interval and high rate under high disease pressure or where conditions are conducive for disease development. DO NOT apply more than 2 applications of OzCrop Cyprofludox WG Fungicide per crop.
	Grey Mould (Botrytis Rot) <i>Botrytis cinerea</i>		Apply OzCrop Cyprofludox WG Fungicide prior to or at the onset of disease. Best results are achieved using a preventative control program, with the first spray commencing soon after transplanting on a 7 to 14 spray interval. Use a 7 day spray interval and high rate under high disease pressure or where conditions are conducive for disease development. DO NOT apply more than 2 applications of OzCrop Cyprofludox WG Fungicide per crop.
	Anthracnose (<i>Microdochium panattonianum</i>)		1 kg/ha in a spray volume of 250 – 500 L water/ha
Nursery stock (non-food) and ornamentals (non-food) including seedlings, plugs, potted colour, trees, shrubs, foliage plants, palms, grasses and fruit trees (non-bearing)	Anthracnose (<i>Colletotrichum</i> spp.)	80 g/100 L or 800 g/ha	Apply via calibrated boom or handheld sprayer when conditions favour disease development, prior to or at the first symptoms. DO NOT apply more than 4 applications per crop at 10 to 14 day intervals. DO NOT apply more than 2 consecutive applications before applying at least the same number of sprays from a different mode of action group.
	Grey mould, Rhizoctonia rot (<i>Rhizoctonia</i> spp), Sclerotinia rot (<i>Sclerotinia</i> spp)		Crop Damage This product has demonstrated toxicity under certain situations, application methods, applications rates and weather conditions to nursery stocks including Impatiens, New Guinea Impatiens, Geraniums, Ferns, Anemone, Begonia, Kalanchoe, Saint Paulia and Exacum affine. This product may also leave undesirable spray residue on foliage and flowers. It is recommended that users treat a sample number of plants or a small area prior to whole crop treatment and monitor any phytotoxic effects that will compromise production goals. This action cannot guarantee crop safety as application, environmental and crop conditions may vary from test treatment to whole crop treatment.

CROP	PEST	RATE/Ha	CRITICAL COMMENTS
Onions, Alliums (including bulb onions, spring onions, shallots and garlic)	Neck rot (<i>Botrytis allii</i>)	800 g/ha to 1 kg/ha plus a registered rate of non- ionic surfactant	Apply OzCrop Cyprofludox WG Fungicide at key timings for botrytis control prior to or at the onset of disease by foliar application with boom spray or equivalent. Apply a second application 7 to 14 days after the initial application if conditions continue to remain favourable for disease development. Use a 7 day spray interval and high rate under high disease pressure or when continual wet, cloudy conditions favour for disease development. Apply in sufficient water to ensure thorough and uniform coverage. Applications MUST be made with the addition of a non-ionic surfactant. DO NOT apply more than 2 applications of OzCrop Cyprofludox WG Fungicide per crop per season. To avoid resistance, the product should be rotated with other approved fungicides from different chemical groups.
	Suppression: Black mould <i>Aspergillus niger</i>		
Pyrethrum (<i>Tanacetum cinerariifolium</i>) crops	Winter dieback complex (Ray blight; <i>Phoma ligulicola</i> and Tan Sport <i>Microsphaeropsis tanacetii</i>)	1 kg/ha	Apply in a total spray volume of 400 L/ha. DO NOT apply more than 2 applications per crop per season with an interval of 7 to 14 days between applications.
Strawberries	Crown and Petiole Rots <i>Colletotrichum gloeosporioides</i>	80 g/100L water	Apply OzCrop Cyprofludox WG Fungicide prior to or at the onset of disease. Repeat applications at 7 to 14 day intervals. Use the shorter interval when conditions favour disease development (wet and cloudy conditions). The rate suggested is for high volume dilute spraying to the point of “run-off”. For concentrate spraying, adjust dilution rate accordingly. DO NOT exceed 800 g/ha. Apply in sufficient water to ensure thorough and uniform coverage of foliage and plant crowns. DO NOT apply more than 3 applications of OzCrop Cyprofludox WG Fungicide per crop.
	Grey Mould (<i>Botrytis cinerea</i>)		Commence a program of protectant fungicides during flowering (blossom) when conditions favour disease. Repeat applications of OzCrop Cyprofludox WG Fungicide on a 7 to 10 day interval. Use the shorter interval when conditions favour disease development. The rate suggested is for high volume spraying to the point of “run-off”. For concentrate spraying, adjust dilution rate accordingly. Apply in sufficient water to ensure thorough and uniform coverage of foliage and fruit. Reduce background levels of disease by removing infected plant debris and rotten fruit. This use is subject to a CropLife fungicide resistance management strategy. DO NOT apply more than 3 applications of OzCrop Cyprofludox WG Fungicide per crop (see Resistance Management).

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

HARVEST:

Alliums (Bulb onions, Spring Onions, Shallots and Garlic), Capsicum, Lettuce, Leafy Vegetables, Onions: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION

Grapes: DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION

Strawberries: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION

Cut Flowers, Nursery Stock (non-food), Ornamentals (Seedlings, plugs, potted colour, trees, shrubs, foliage plants, palms, grasses and non-bearing fruit trees), Pyrethrum: NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

OzCrop Cyprofludox WG Fungicide is a water dispersible granule prepack mixture of cyprodinil and fludioxonil. Cyprodinil is systemic compound that is taken up into the cuticle and waxy layers of leaves and fruit. From there it is distributed and penetrates into other parts of the plant. Fludioxonil is the residual component of the mixture and stays mainly on the leaf and fruit surfaces.

Application

Ground Application Only: Ensure thorough coverage of foliage and fruit (or bunches) using equipment delivering a MEDIUM spray quality. Ensure that the correct amount of OzCrop Cyprofludox WG Fungicide is applied per hectare irrespective of water application rate per hectare.

Dilute Spraying

Use a sprayer designed to apply high volumes of water up to the point of runoff and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of runoff. Avoid excessive runoff. The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice. Add the amount of product specified in the Direction for Use table for each 100 L of water. Spray to the point of runoff. The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

Concentrate Spraying

Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of runoff) and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume. Determine an appropriate dilute spray volume (see Dilute Spraying) for the crop canopy. This is needed to calculate the concentrate mixing rate. The mixing rate for concentrate spraying can then be calculated in the following way:

Example Only

1. Dilute spray volume as determined above: for example 1000 L/ha
2. Your chosen concentrate spray volume: for example 400 L/ha
3. The concentration factor in this example is: $2.5 \times$ (ie $1000 \text{ L} \div 400 \text{ L} = 2.5$)
4. If the dilute label rate is 80 g/100 L, then the concentrate rate becomes 2.5×80 , that is 200 g/100 L of concentrate spray.

The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to be changed as the crop grows. For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

Mixing

OzCrop Cyprofludox WG Fungicide is a water dispersible granule (WG) fungicide that mixes readily with water. Partly fill the spray tank with water. Start the agitation and add the correct amount of product to the spray tank with the agitation system running. Continue agitation while topping up the spray tank with water. Add a non-ionic surfactant at registered rates (where required) at the completion of filling of the spray tank. Continue agitation while spraying.

Tank Mixing:

When mixing OzCrop Cyprofludox WG Fungicide and other water dispersible granule (WG) or wettable powder (WP) formulations, ensure they are added and mixed well prior to adding emulsifiable concentrate (EC) or suspension concentrate (SC) products. Wettable powder (WP) formulations should be pre-mixed separately and then added to the spray tank.

Compatibility

OzCrop Cyprofludox WG Fungicide is compatible with Bugmaster* Flowable, Delfin* WG, Dithane*, Gusathion*, Kelthane*, Kocide*, Lorsban* 500 W, Polyram DF, RIDOMIL®GOLD PLUS, Sanmite*, SUPRACIDE®, THIOVIT JET®, and TOPAS®. Mixtures with more than one of the above are not recommended. As formulations of other manufacturer's products are beyond the control of OZCROP Pty Ltd, all mixtures should be tested prior to mixing commercial quantities. OzCrop Cyprofludox WG Fungicide may not be compatible with Bravo®.

Fungicide Resistance Warning

OzCrop Cyprofludox WG Fungicide is a combination of an anilinopyrimidine and a phenylpyrrole fungicide. For fungicide resistance management, OzCrop Cyprofludox WG Fungicide is both a Group 9 and a Group 12 fungicide. Some naturally occurring individual fungi resistant to OzCrop Cyprofludox WG Fungicide and other Group 9 and Group 12 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group 9 and Group 12 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use Shandong Rainbow International Co Ltd accepts no liability for any losses that may result from the failure of this product to control resistant fungi.

GROUP	9	12	FUNGICIDE
-------	----------	-----------	-----------

Resistance Management

In order to minimise the likelihood of the development of fungicide resistance, it is recommended that OzCrop Cyprofludox WG Fungicide be used in a program incorporating all available forms of crop protection. OzCrop Cyprofludox WG Fungicide should be applied as part of a protective spray program using fungicides from different mode of action group/s. Alternating or mixing fungicides with different modes of actions is a recognised disease resistance management strategy. DO NOT wait until disease levels have built up before commencing applications as this reduces the effectiveness of control and increases the risk of resistance development. Disease control may be reduced if strains of pathogens less sensitive to OzCrop Cyprofludox WG Fungicide develop. OzCrop Cyprofludox WG Fungicide should be applied as specified in the Directions For Use in association with the following CropLife Fungicide

Resistance Management Strategies:

Grapes: If 3 or fewer bunch rot sprays are applied in a season, use only 1 spray containing a Group 9 fungicide. If 4 or more bunch rot sprays are applied in a season, use a maximum of 2 sprays containing a Group 9 fungicide. Note: In the absence of an established resistance management strategy for a particular crop/disease situation, it is recommended that:

The use of fungicides from any given activity group (excluding Group M) be limited to a maximum of 1/3 of the total number of fungicide applications per crop. A maximum of 2 consecutive applications of OzCrop Cyprofludox WG Fungicide are to be applied, including from one season/crop to the next.

Capsicums and Strawberries: If 3 or fewer Botrytis fungicide sprays are applied per crop, use only 1 spray containing a Group 9 fungicide. If 4 to 6 sprays are applied per crop use a maximum of 2 sprays containing a Group 9 fungicide. If 7 or more sprays are applied per crop use a maximum of 3 sprays containing a Group 9 fungicide. Note: In the absence of an established resistance management strategy for a particular crop/disease situation, it is recommended that:

The use of fungicides from any given activity group (excluding Group M) be limited to a maximum of 1/3 of the total number of fungicide applications per crop.

A maximum of 2 consecutive applications of OzCrop Cyprofludox WG Fungicide are to be applied, including from one season/crop to the next.

Export Wine

Where this product will be used on grapes that will be used to produce wine destined for export markets, seek advice from the Australian Wine Research Institute to ensure the wine will meet the requirements of the intended importing country.

PRECAUTIONS

Re-entry Period:

DO NOT enter treated area until the spray has dried unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each days use.

INTEGRATED PEST MANAGEMENT

This product can be used where integrated pest management is being practiced.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers. DO NOT apply if heavy rains are imminent. DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto adjacent areas, particularly wetlands, water bodies, or watercourses.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, dry, well ventilated area, out of direct sunlight. Single rinse liner before disposal. Add rinsings to spray tank.

[hermetically sealed bag]

Single rinse liner before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. Puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

[HDPE container]

Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. May irritate the nose and throat. Will irritate the eyes. Repeated exposure may cause allergic disorders. Avoid contact with eyes. Do not inhale dust or spray mist. When opening the container and preparing spray or dip, and using the prepared spray or dip, wear cotton overalls buttoned to the neck and wrist and elbow length chemical resistant gloves and face shield or goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

Safe Work Australia:

May cause an allergic skin reaction.
May cause damage to organs through prolonged or repeated exposure.
Very toxic to aquatic life with long lasting effects.
Do not breathe dusts.
Do not get in eyes, on skin, or on clothing.
Avoid release to the environment
Get medical advice or attention if you feel unwell.
Brush off loose particles from skin.
IF SWALLOWED: Rinse mouth. Do NOT induce vomiting.
IF ON SKIN: Wash with plenty of soap and water.
If skin irritation or rash occurs: Get medical advice.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet (SDS), which can be obtained from the supplier.

CONDITIONS OF SALE

The use of OzCrop Cyprofludox WG Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by OzCrop Pty Ltd, regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and OzCrop Pty Ltd accepts no responsibility for any consequence whatsoever from the use of this product.



OzCrop Pty Ltd ABN 16 160 656 431

G13/25 Solent Circuit, NORWEST NSW 2153

Telephone: (02) 8123 0170 • Facsimile: (02) 8123 0171

Email: enquiries@ozcrop.com.au

visit our website www.ozcrop.com.au

APVMA Approval No.: 83349/124103

Label Release: 202103

