

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

OzCrop

Diuron 900 WG

HERBICIDE

ACTIVE CONSTITUENT: 900 g/kg DIURON

GROUP **C** HERBICIDE

For selective weed control on certain crops as shown under the Directions for Use

PACK SIZES:

15 kg

**NOT A DANGEROUS GOOD
ACCORDING TO THE
AUSTRALIAN DANGEROUS
GOODS (ADG) CODE.**

OzCrop Pty. Ltd.
ABN 16 160 656 431
Suite G13/25 Solent Circuit,
Norwest, NSW 2153
TEL: 02 8123 0170

DIRECTIONS FOR USE:**RESTRAINTS**

DO NOT use in water logged areas

DO NOT apply if greater than 50 mm rainfall is expected within 3 days of application

DO NOT irrigate within 3 days of irrigation

DO NOT apply to fields where the slope exceeds 3%

DO NOT spot spray more than 5% of total farm areas

DO NOT apply more than once per calendar year

DO NOT apply when wind speed is less than 3 or more than 20 kilometres per hour as measured at the application site

Cotton

DO NOT use in cotton UNLESS ALL irrigation tailwater and/or up to 25 mm rainfall can be captured and held on farm

Sugar Cane

Post-emergent use (1.8 kg ac/ha max 60% band spray)

APPLY ONLY as a directed band spray over a maximum of 60% of the crop area

DO NOT apply in the Wet Tropics

DO NOT apply in the Burdekin between 1 January and 29 February

DO NOT apply in Mackay/Whitsunday between 1 December and 30 April

DO NOT apply in Mary-Burnett between 1 November and 29 February

DO NOT apply in NSW between 1 November and 30 April

Irrigation channels and drains

DO NOT use in irrigation channels or drains UNLESS ALL irrigation tailwater and rainfall can be captured and held on farm

Desert channels (bore drains)

DO NOT apply between 1 December and 30 March

DO NOT open drains for 72 hours following treatment

SPRAY DRIFT RESTRAINTS

DO NOT apply by air

DO NOT apply with spray droplets smaller than COARSE spray droplet size category according to nozzle manufacturer specifications that refer to the ASAE S572 Standard or BCPC Guideline

Users of this product MUST make an accurate written record of the details of each spray application within 24 hours following application and KEEP this record for a minimum of 2 years. The spray application that must be recorded are: 1. Data with start and finish times of application; 2. Location address and paddock/s sprayed; 3. Full name of this product; 4. Amount of products used per hectare and number of hectares applied to; 5. Crop/situation and weed/pest; 6. Wind speed and direction during application; 7. Air temperature and relative humidity during application; 8. Nozzle brand, type, spray angle, nozzle capacity and spray system pressure during application; 9. Name and address of person applying this product. (Additional record details may be required by the State or Territory where this product is used).

MANDATORY NO SPRAY ZONES

DO NOT apply when there is non-target vegetation downwind from the application area and within the mandatory no-spray zones shown in table below

DO NOT apply when there are aquatic and wetland areas including aquacultural ponds, surface streams and rivers downwind from the application area and within the mandatory no-spray zones shown in table below.

SITUATION	RATE (g ac/ha)	DOWNWIND BUFFER ZONE (m)	
		AQUATIC	TERRESTRIAL
Wheat, barley, triticale, cereal rye, oats (WA only)	250-500	30	60
Wheat, barley, triticale and oats (WA only)	180-250	15	30
Wheat, barley and oats (NSW, Vic, ACT, SA only)	450	25	50
Wheat and barley – SA only	640-880	50	100
Wheat and barley – NSW, Vic, ACT, SA only	250	15	30
Summer fallows	250	15	30
Cotton	900-1800	100	200
Lupins – WA only	990	30	80
Sugarcane, bananas	250-450	25	50
Sugarcane	1800	100	200
Asparagus	1800	100	200

CROP	WEED	STATE	RATE/ha	CRITICAL COMMENTS
Asparagus	Annual broadleaf weeds and some annual grasses	VIC only	1.5-2 kg	Apply in established beds after they have been "hilled up" and before emergence of spikes. This will give 10-14 weeks freedom from weeds. Post-emergence application may be made, but the spray should be directed so as to contact only the base of the stems. Use higher rates on heavier soils. DO NOT use on seedling asparagus.
Bananas	Amaranthus, Barley Grass, Barnyard Grass, Capeweed, Cobbler's Pegs, Crowsfoot Grass, Fat Hen, Guinea Grass, Mustard, Pigweed, Prince of Wales Feather, Rye Grass, Sow Thistle, Summer Grass, Wild Oats, Wild Radish, Wild Turnip	Qld, NSW, ACT only	275-500 g Plus 1.2 or 1.6 L/ha OzCrop Paraquat 250	Add to OzCrop Paraquat 250 to enhance weed control under favourable growing conditions and in open sunny conditions. DIURON 900 WG at rates up to 500 g/ha can be blanket sprayed. Use a directed spray for higher rates of DIURON 900 WG. Complete spray coverage is essential. For grasses and broadleaved weeds up to 5 cm high use a minimum of 250 L spray solution/ha, increase to 350 L/ha for weeds up to 10 cm high. Use a minimum spray volume of 400 L/ha for weeds > 10 cm high and for dense mature weeds. Refer to OzCrop Paraquat 250 label for rates and full directions. If OzCrop Paraquat 250 rate is less than the ratio 500 mL/100 L add 100 mL non-ionic surfactant 600 g/L or 60 mL non-ionic surfactant 1000 g/100 L spray mix.
Cotton	Annual broadleaf weeds and some annual grasses	Qld, NSW, ACT only	1-2 kg	Pre-emergence: Apply at planting or within 7 days of planting and prior to cotton emergence. Seed should be planted at least 4 to 5 cm deep and soil compacted over seeds by a normal planter packer wheel. Use high rate on soils high in clay or organic matter content. DO NOT use on light sandy soils as injury to crop may occur. DO NOT make a second pre-emergent application to the same field during the same crop year as injury to the crop may result. Apply in 300 to 450 L water/ha. Post-emergence (lay-by): Use the higher rate on heavier soil types high in clay and/or organic matter. DO NOT use on light sandy soils low in organic matter as injury to the cotton may result. Apply when cotton is at least 30 cm high, or immediately after the last cultivation. If emerged annual weeds (15 cm or less in height) are present add 170 mL non-ionic surfactant 600 g/100 L water to the spray solution. Apply DIURON 900 WG as a directed spray to cover the area beneath cotton plants and between rows. Avoid spray contact or drift on to cotton plants as injury may result. DO NOT apply more than once per season. DO NOT re-plant treated areas to any crop within one year after last application as injury may result except cotton, corn or sorghum (grain sorghum only) which may be planted in the Spring following treatment. Apply 300-500 L water/ha.
Lupins (on light soils only)	Annual Ryegrass, Capeweed, Clovers, Doublegee, Medics, Wild Radish, Wild Turnip and suppression of Dock, Sorrel, Wireweed, Wild Oats and Brome Grass	WA only	1.1 kg	Use through boom spray equipment only. Apply in a minimum of 150 L water/ha. Higher volumes of water will give more even application. Must be applied prior to crop emergence. DO NOT use on white or grey sands.

CROP	WEED	STATE	RATE/ha	CRITICAL COMMENTS
Sugar cane	Amaranthus, Barley Grass, Barnyard Grass, Capeweed, Cobbler's Pegs, Crowsfoot Grass, Fat Hen, Guinea Grass, Mustard, Pigweed, Prince of Wales Feather, Ryegrass, Sowthistle, Summer Grass, Wild Oats, Wild Radish, Wild Turnip	Qld, NSW only	1.9 kg	Pre-emergence of cane and weeds. Soil surface must be moist at all times of application. Rain or irrigation is required within 10 days of application.
			1.9 kg	Post-emergence of cane, as for pre-emergence. Use lower rate if following an application at planting. If weeds at early post-emergence stage addition of non-ionic surfactant 600 g/L at 250 mL/100 L of water will enhance activity, but direct spray to below cane leaves.
			275 -1.9 kg plus 1.2-1.6 OzCrop Paraquat 250	Add to OzCrop Paraquat 250 to enhance weed control under favourable growing conditions and in open sunny conditions. DIURON 900 WG at rates up to 500 g/ha can be blanket sprayed. Use a directed spray for higher rates of DIURON 900 WG. Complete spray coverage is essential. For grasses and broadleaved weeds up to 5 cm high use a minimum of 250 L spray solution/ha. Increase to 350 L/ha for weeds up to 10 cm high. Use a minimum spray volume of 400 L/ha for weeds > 10 cm high and for dense mature weeds. Refer to OzCrop Paraquat 250 label for rates and full directions. Note: Where broadleaved weeds are a serious problem add Tornado* DF to the mixture.
Summer Fallow	Potato Weed (<i>Heliotropium europaeum</i>)	SA only	275 g	Only effective when mixed with OzCrop Blowout (diquat/paraquat) at 1.6 L/ha (See OzCrop Blowout label).
Wheat, Barley, Triticale, Cereal Rye, Oats	Capeweed, Doublegee, Mustard, Wild Radish (suppression only), Water Weeds (Toad Rush, Crassula)	WA only	275-550 g *Barley to 390 g only	Post planting application only for full cut sowing systems that leave a relatively smooth surface, eg. standard combine or airseeder. Apply post plant pre-emergent to crop within 3-4 days of planting.
	As above plus suppression of Annual Ryegrass, Barley Grass and Silver Grass		550 g plus 500 mL of a 720 g/L metolachlor to improve suppression of Ryegrass	Pre plant application only for sowing systems (eg. No Till or Zero Till) which leave open slots or furrows, or where seed is sown along the bottom of the furrow. Even the soil surface where harrowed after sowing. Apply within 7-10 days pre plant. Soil Types – use lower rates on sandy surfaced soils with a low organic carbon or clay content. Caution: DO NOT use on grey or white sands. Crop to be planted at least 35 mm deep. Crop tipping and yellowing may occur particularly when heavy rain falls after sowing. Shallow sown crops are more at risk. Caution: Severe crop damage will occur if herbicides containing Paraquat or Diquat contact an emerging cereal crop either as a salvage treatment or via spray drift unless rainfall has been exceptionally low.
Wheat, Barley, Triticale, Cereal Rye			275-550 g plus 830 mL/ha of trifluralin 480 g/L	As above. Must be incorporated within 4 hours of application.

CROP	WEED	STATE	RATE/ha	CRITICAL COMMENTS
Wheat, Barley, Oats	Amsinkia (Yellow Burr), Capeweed, Charlock, Deadnettle, Iceplant, Melilotus, Mustards, Poppies, Saffron Thistle, Sheepweed, Spiny Emex, Three Corner Jack, Wild Radish, Wild Turnip	NSW, ACT, Vic, SA only	500 g	Apply when weeds are at the 2-4 leaf stage and soon after the crop is past the 2-5 leaf stage which is normally within 6 weeks of sowing. DO NOT spray later than early tillering stage.
Wheat, Barley, Triticale, Oats	Capeweed, Doublegee, Erodium, London Rocket, Lupin, Mustard, Rapistrum, Wild Radish, Wild Turnip, Storksbill, Wild Mustard	WA only	200 g plus 400 mL MCPA 500	Apply when the crop has at least 3 to 5 leaves on the main stem (Zadoks 13-15). Apply by boom spray. Weeds must be in the 2-5 leaf stage. Crop and weeds should be dry at the time of spraying. DO NOT spray if rain imminent.
			280 g plus 200 mL 2,4-D Amine 625	Apply when the crop has at least 4 to 5 leaves on main stem (Zadoks 14-15). Undersown sub-clovers may be slighted retarded. DO NOT apply to undersown medics. Apply in 40-70 L water/ha.
Wheat and Barley	Soursob	SA only	710 g-980 g	Boom Spray: Sow the crop at least 5 cm deep. Apply when crop is in the 2-5 true leaf stage. Use the lower rate on lighter soils.
	Amsinkia (Yellow Burr), Capeweed, Deadnettle, Mustards, Doublegee (Three Corner Jack), Iceplant, Mustards, Turnips, White Ironweed (Sheepweed), Wild Radish	NSW, ACT, Vic only	280 g plus 500 mL MCPA 500	Apply by boom spray when the crop is at the 3-5 leaf stage (NSW, SA) or 5 leaf stage (Vic). Weeds must be at the 2-5 leaf stage. DO NOT spray if rain is imminent. DO NOT apply to undersown medics or lucerne. DO NOT use on Durati, Hybrid Titan, Kite, Kamilaroi or Shortim Wheat.
SA only		280 g plus 350 mL MCPA 500		
Irrigation channels and drainage ditches	Annual grass, broadleaf weeds and perennial grasses (except Bindweed, Nutgrass and Russian Knapweed)	Qld, Vic, Tas, SA, WA only	22 kg or 220 g per 100/m ²	High Volume or Power Spraying: Apply in 2000 L water. Apply during the non-crop season and when channel is not in use. Soil should be moist and preferably bare i.e. after regrading and cleaning channels. Apply prior to expected seasonal rainfall so as to "fix" it into the soil. If 50-100 mm of rain has not fallen after application and it is necessary to use the irrigation channel, fill it with water and allow to stand for 72 hours, then drain off the water in the channel and run into wasteland.
		NSW only	40 kg maximum	
Bore Drains	Mimosa Bush, Prickly Acacia	Qld only	35.5 kg	Block off bore drain 24 hours prior to spraying and allow drain to empty. Spray 1 m wide strip onto mud. Wait 48-72 hours before opening drain again. Water MUST by-pass house tanks and desirable tree stands for 7 days after opening drain. DO NOT delve for 2 weeks.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

DO NOT ALLOW ANIMALS TO DRINK WATER FROM TREATED BORE DRAINS FOR 3 DAYS BEFORE SLAUGHTER FOR HUMAN CONSUMPTION.

GRAZING / HARVEST:

NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

Before opening, carefully read Directions For Use, Precautionary Statements, Safety Directions and First Aid Instructions.

Application should be made to moist soil before weeds emerge unless otherwise recommended. Sufficient moisture in the form of rain or irrigation is necessary after application to move the chemical into the weed root zone. Add non-ionic surfactant where recommended. When spraying emerged weeds best results are obtained when weeds are actively growing.

Thoroughly clean all traces of this product from application equipment immediately after use, otherwise injury to desirable vegetation may result when equipment is used again. Flush tank, pump, hose and boom with several changes of water after removing nozzle tip and screens (clean these parts separately). Keep from contact with fertilisers, insecticides, fungicides and seeds.

RESISTANT WEED WARNING

OzCrop DIURON 900 WG Herbicide is a

GROUP	C	HERBICIDE
-------	----------	-----------

 member of the Urea group of herbicides. DIURON 900 WG has the Inhibitor of photosynthesis at photosystem II mode of action. For weed resistance management DIURON 900 WG is a Group C herbicide.

Some naturally-occurring weed biotypes resistant to DIURON 900 WG and other Group C herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by DIURON 900 WG or other Group C herbicides.

However resistance should not be assumed without first viewing the method of application, the coverage and the timing of application. Resistance to DIURON 900 WG cannot be predicted and may occur at any time. It is not due to a fault in the product and OZCROP Pty. Ltd. accepts no liability for any losses that may result from the failure of DIURON 900 WG to control resistant weeds.

MIXING

Part fill tank with CLEAN water (50%). Commence vigorous agitation. Add granules steadily. DO NOT dump. Ensure complete dispersion before filling to volume with water. Continue agitation during spraying. DO NOT leave spray solution in tank overnight.

Agitation: An efficient by-pass agitator or means of mechanical agitation should be employed when using this product.

COMPATIBILITY

This product is compatible with diquat, OzCrop Paraquat 250 SL, amitrole, 2,4-D Amine and Ester, 2,4-DB, MCPA, chlorpyrifos, omethoate and chlorsulfuron. Add the required amount of product when the spray tank containing DIURON is nearly full and being continually agitated. This product is not compatible with methabenzthiazuron, dimethoate, flamprop-m-methyl, OzCrop Diclofop-Methyl 375 and triadimefon.

As formulations of other manufacturers' products are beyond the control of OZCROP, all mixtures should be tested prior to mixing commercial quantities.

WEED CONTROL ON NON-CROP LAND

DIURON 900 WG is an effective herbicide for the residual control of many annual and perennial grasses and broadleaf weeds on non-crop land such as irrigation and drainage ditches where bare ground is desired. Established perennials such as Bindweed, Johnson Grass, Nutgrass and Russian Knapweed are not satisfactorily controlled at rates recommended. Duration of control depends on the amount of chemical applied, soil type, rainfall and particular weed species.

SELECTIVE USE IN CROPS

For selective use in crops it is important to apply recommended rates correctly. DIURON 900 WG is most effective when applied to soil which has been well tilled (clod free) and is free of weeds and trash. Where not recommended for overall application, care should be taken to avoid spray drift onto fruit or foliage. Wetting agents should only be used where specified.

REPLANTING RESTRICTIONS

Crop Rotations: DO NOT replant treated areas to any crop within two years after last spray unless otherwise stated.

Cotton – DO NOT replant treated areas to any crop within 1 year after last spray. Except cotton, corn or grain sorghum which may be planted in the spring of the following year.

Sugar Cane – Treated areas can be replanted to sugar cane 1 year after last spray.

CROP SAFETY

Heavy rains after application of this product may cause severe crop damage, risk must be assumed by user.

DO NOT use on sandy or gravelly soils or soils low in organic matter. Avoid contact of sprays with crop foliage or fruit.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied at any time as recommended in the Directions For Use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used container.

STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN. Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Shake empty bag into spray tank. DO NOT dispose of undiluted chemicals on-site. Puncture or shred and bury empty containers at a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Avoid contact with eyes and skin. DO NOT inhale dust or spray mist. If product on skin immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

MSDS

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for DIURON 900 WG is available from OZCROP on request. Call Customer Service on (02) 8199 3610.

CONDITIONS OF SALE: The use of OzCrop DIURON 900 WG Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by OZCROP Pty. Ltd., regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and OZCROP Pty. Ltd. accepts no responsibility for any consequence whatsoever from the use of this product.

* Other trademarks

Safe Work Australia:
Harmful if swallowed.
Suspected of causing cancer.
May cause damage to organs through prolonged or repeated exposure.
Very toxic to aquatic life with long lasting effects.
Do not eat, drink or smoke when using this product.
Use personal protective equipment as required.
Wash contaminated clothing before reuse.
IF SWALLOWED: Rinse mouth. Do NOT induce vomiting.