

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

 OzCrop

Haloxyfop 520 EC

HERBICIDE

ACTIVE CONSTITUENT: 520 g/L HALOXYFOP
present as the HALOXYFOP-R-METHYL ESTER

GROUP A HERBICIDE

For the post-emergent control of a wide range of annual and perennial grass weeds in grain legume and oilseed crops, lucerne, medic and clover pasture and seed crops, forestry, bananas, citrus, grapes, pineapples, pome and stone fruit, pyrethrum, tropical fruit and nut crops as specified in the Directions for Use.

CONTENTS:

5L, 10L, 20L

**NOT A DANGEROUS GOOD
ACCORDING TO THE AUSTRALIAN
DANGEROUS GOODS (ADG) CODE.**

BATCH NO.

DATE OF MANUFACTURE:

OzCrop Pty Ltd ABN 16 160 656 431
Suite G13/25 Solent Circuit, Norwest, NSW 2153
Tel: 02 8123 0170 • Fax: 02 8123 0171
Email: enquiries@ozcrop.com.au
www.ozcrop.com.au

DIRECTIONS FOR USE:

Restraint: DO NOT apply to weeds which may be stressed (not actively growing) due to prolonged periods of extreme cold, moisture stress (waterlogged or drought affected), poor nutrition or previous herbicide treatment as reduced levels of control may result.
DO NOT spray if rain is likely to occur within one hour.

Table 1a. Winter crops – Canola, Chickpeas, Faba beans, Field peas, Lentils, Linola, Linseed, Lupins, Lucerne, Vetch, Medic and Clover pastures or seed crops.

WEEDS CONTROLLED	WEED GROWTH STAGE	Rate (mL/ha)		CRITICAL COMMENTS
		With Uptake ¹ Spraying Oil	With non-ionic wetter ²	
Annual Ryegrass	2 to 4 leaf	75	100	^{1,2} See GENERAL INSRUCTIONS , Spraying oils/wetters section. CANOLA, LINOLA AND LINSEED: DO NOT apply after the 8 leaf stage of the crop. DO NOT apply after the commencement of stem elongation. This means that application must not occur after the 8 leaf stage, or if stem elongation commences before the 8 leaf stage, application must not occur after stem elongation has commenced. DO NOT apply more than 1 application of herbicide containing haloxyfop per crop. DO NOT apply after grazing.
	Early tillering	100	100	
Barley Grass	2 to 4 leaf	50	75	FIELD PEAS AND CANOLA: The only oil recommended for use with OzCrop Haloxyfop 520 EC Herbicide is Uptake* Spraying Oil. OzCrop Haloxyfop 520 EC Herbicide + Lontrel 750 SG + Uptake* Spraying Oil are compatible and selective to canola. This tank-mix is also compatible with atrazine and selective to triazine tolerant canola. LUPINS AND FIELD PEAS: Mixtures with Brodal or simazine may cause crop yellowing and separate applications are recommended. CHICKPEAS, FABA BEANS, LENTILS AND VETCH, LINOLA, LINSEED: Broadleaf herbicides should not be added to OzCrop Haloxyfop 520 EC. Apply OzCrop Haloxyfop 520 EC Herbicide and broadleaf herbicides at least a week apart. LUCERNE, CLOVER OR MEDIC PASTURES: If grazed or cut for hay immediately prior to treatment delay application until all grasses have fully expanded leaves. Use 75 mL + spraying oil or 100 mL + wetter/ha. (See GENERAL INSTRUCTIONS , Spraying Oils/wetters section). If Silver Grass (<i>Vulpia</i> spp.) is present in pasture, simazine should be tank mixed with the higher rate of OzCrop Haloxyfop 520 EC Herbicide plus a non-ionic wetter.
Brome Grass	Early tillering	75	100	
Paradoxa Grass				
Volunteer Cereals				
Wild Oats	2 to 4 leaf	37.5	50	
WA, SA, Vic, Tas, southern & central NSW	Early tillering	50	75	
Wild Oats	2 to 4 leaf	50	75	
northern NSW & Qld	Early tillering	75	100	

Table 1b. Winter crop growth stage application windows.

CROP	CROP GROWTH STAGE
Lucerne, Medic and Clover pastures or seed crops.	Apply from 2nd trifoliate leaf onwards. For <i>Erodium</i> spp. spraying, apply from cotyledon crop stage onwards.
Canola, Linola and Linseed	Apply from 2 leaf to 8 leaf stage of crop growth. DO NOT apply after the commencement of stem elongation. This means that application must not occur after the 8 leaf stage, or if stem elongation commences before the 8 leaf stage, application must not occur after stem elongation has commenced.
Chickpeas, Faba beans, Field peas, Lentils, Lupins, Vetch	Apply from 2nd leaf, 2nd node or 2nd branch prior to flowering.

Table 2a. Lucerne, Medic and Clover seed crops and pastures. See table 1b for crop stages.

WEEDS CONTROLLED	WEED GROWTH STAGE	RATE (mL/ha) with Uptake ¹ Spraying Oil	CRITICAL COMMENTS
Prairie Grass (<i>Bromus catharticus</i>)	Up to early tillering	100	¹ See GENERAL INSRUCTIONS , Spraying oils/wetters section. ³ Use lower when growing conditions and crop or pastures competition are good and when weed populations are below 100 plants/m ² . Use the higher rate when weed populations are above 100 plants/m ² or when crop or pasture competition is poor. Note: Storksbill may not be controlled if simazine or Broadstrike are tank-mixed with OzCrop Haloxyfop 520 EC Herbicide.
Musky or Ferny Leaf Storksbill (<i>Erodium moschatum</i>)	Up to 6 leaf or 5 cm diameter	50 – 75 ³	LUCERNE, CLOVER OR MEDIC PASTURES: If grazed or cut for hay immediately prior to treatment delay application until all grasses have fully expanded leaves. Use 75mL + spraying oil or 100 mL + wetter/ha. (See GENERAL INSTRUCTIONS , Spraying oils/wetter section). If Silver Grass (<i>Vulpia</i> spp.) is presented in pasture, simazine should be tank mixed with the higher rate of OzCrop Haloxyfop 520 EC Herbicide plus a non-ionic wetter.
Common Crowsfoot or Common Storksbill (<i>Erodium cicutarium</i>)			
Long or Shiny Leaf Storksbill (<i>E. botrys</i>)	Up to 8 leaf or 5 cm diameter	75 – 100	

Table 2b. Lucerne, Medic and Clover seed crops only – not to be used for stockfeed. See table 1b for crop stages.

WEEDS CONTROLLED	WEED GROWTH STAGE	RATE (mL/ha) with Uptake ¹ Spraying Oil	CRITICAL COMMENTS
Couch Grass (suppression)	Tillering seedlings	150 + 150 ⁴	¹ See GENERAL INSRUCTIONS , Spraying oils/wetters section. ⁴ For best suppression of couch or control of Rhodes Grass, make 2 applications of OzCrop Haloxyfop 520 EC Herbicide 2-4 weeks apart. Time second application to coincide with tillering stage of weeds and just after irrigation or significant rain.
Rhodes Grass (control)			Only treat actively growing weeds, which are not moisture stressed. Use these rates for control of Couch and Rhodes Grass.
Couch Grass (control)	Established stands	400 – 800	
Rhodes Grass (control)			

Table 3a. Summer Crops – Cotton, Cowpea, Lucerne, Mung bean, Navy beans, Peanuts, Soybeans, Sunflowers

WEEDS CONTROLLED	WEED GROWTH STAGE	RATE (mL/ha) with Uptake ¹ Spraying Oil	CRITICAL COMMENTS
Australian Millet	2 leaf to tillering up to 15 cm	150	¹ See GENERAL INSRUCTIONS , Spraying oils/wetters section. NAVY BEAN, PEANUTS, SOYBEANS For broadleaf weed control, OzCrop Haloxyfop 520 EC Herbicide at 150 mL/ha plus wetter may be tank mixed with Blazer* (except on Navy beans) or Basagran*. Tank mixes may cause transient leaf spotting on the crop but do not normally affect yield. DO NOT tank mix broadleaf herbicides with OzCrop Haloxyfop 520 EC Herbicide if grasses have begun tillering or if the grasses are under moisture stress. DO NOT add Uptake Spraying Oil with Blazer or Basagran. DO NOT use Blazer or Basagran tank-mixes on Cowpeas.
Barnyard Grass	2 to 5 leaf	100	
	Tillering up to 15 cm	150	
Crowsfoot Grass	2 leaf to tillering up to 15 cm	150	
Green Panic Grass			
Johnson Grass (Rhizome)			
Johnson Grass (seedling)	2 to 5 leaf	100	
Liverseed Grass (seedling)	Tillering up to 15 cm	150	
Mossman River Grass			
Summer Grass	2 leaf to tillering up to 15 cm	150	
Volunteer Cereals	2 to 4 leaf	100	
	Tillering up to 15 cm	150	



Table 3b. Summer crop growth stage application windows.

CROP	CROP GROWTH STAGE
Lucerne	Apply from 2nd trifoliate leaf onwards.
Cowpea, Mung beans, Navy beans, Soybeans	Apply from 2nd leaf to flowering.
Peanuts	Apply from 5cm to pegging.
Cotton	Apply from 2nd leaf to before the onset of flowering.
Sunflowers	Apply from 2nd leaf to head initiation.

Table 4. Annual and Perennial grasses and *Erodium* spp. In Orchard, Vine, and Plantation crops, Forestry and Pyrethrum.

CROP	CROP GROWTH STAGE	WEEDS CONTROLLED	WEED STAGE	Rate (mL/ha) with Uptake ¹ Spraying Oil	CRITICAL COMMENTS
Orchard, Vine and plantation Crops including: Apples, Avocado, Banana, Blueberry, Citrus, Custard apple, Feijoa, Grapevines, Guava, Kiwifruit, Litchi (Lychee), Longan, Mango, Nashi, Nut trees, Passionfruit, Paw paw, Pear, Persimmon, Pineapple, Rambutan, Stone Fruit	All growth stages	Perennial Grasses, Couch, Rhodes Grass, Slender Rats Tail Grass	Established stands	400 – 800	¹ See GENERAL INSTRUCTIONS , Spraying oils/wetters section.
		Buffel Grass, Green Panic, Johnson Grass, Kikuyu, <i>Paspalum</i> spp., <i>Setaria</i> spp.	Vegetative to early tillering	200	Spray should be directed to the base of the tree or vine avoiding contact with fruit and foliage.
		Annual Grasses: Annual Ryegrass, Barley Grass, Barnyard Grass, Brome Grass, Crowsfoot Grass, Lesser Canary Grass, Liverseed Grass, Mossman River Grass, Paradoxa Grass, Summer Grass, Volunteer cereals, Wild Oats	Late tillering	400	Spot spray: Use 25 mL to 50 mL/100 L of water. Use higher rate on late tillering mature grasses.
Forestry: <i>Pinus radiata</i> , <i>Eucalyptus</i> spp.			2 leaf to tillering	200	Annual Grasses: Where treated in association with perennial grasses, these annual grasses will be controlled
Forestry: <i>Pinus pineaster</i>		Annual grasses as above	Vegetative to tillering	125 – 250	Forestry: For annual grasses apply lowest rate to newly emerged grasses, increasing the rate as they develop.
Pyrethrum		Barley Grass, Brome Grass, Rope Twitch, Barnyard Grass, <i>Erodium</i> spp., Volunteer Cereals	Vegetative to tillering	100 – 250	Pyrethrum (Tasmania only): For <i>Erodium</i> spp. apply 75-100 mL/ha if the main weed is <i>E. botrys</i> . Use 50-75 mL/ha if either <i>E. cicutarium</i> or <i>E. moschatum</i> are the main weeds.

Table 5. OzCrop Haloxypop 520 EC Herbicide and Select Herbicide tank mixes – Canola, Chickpeas, Faba beans, Field peas, Lupins, Lentils

WEEDS CONTROLLED	WEED GROWTH STAGE	Rate (mL/ha)		CRITICAL COMMENTS
		OzCrop Haloxypop 520 EC	Select* Herbicide	
FOP/DIM susceptible Annual Ryegrass + Volunteer Barley, Volunteer Wheat, Brome Grass, Wild Oats, Barley Grass, Phalaris	2 to 4 leaf	25	150	¹ See GENERAL INSTRUCTIONS , Spraying oils/wetters section. Use Uptake Spraying Oil at 500 mL/100 L or Hasten at 1 L/100 L. Apply at the same crop growth stages as those in Table 1b Winter Crops.
	Early tillering	38	150	
FOP resistant Annual Ryegrass + Volunteer Barley, Volunteer Wheat, Brome Grass, Wild Oats, Barley Grass, Phalaris	2 to 4 leaf	25	200	Lentils: Apply up to 7 node-early branching crop growth stage. Lupins: Not for Qld.
	Early tillering	38	250	

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

HARVESTING WITHHOLDING PERIODS

NOT REQUIRED WHEN USED AS DIRECTED FOR:

Orchard Crops, Plantation Crops and Vine Crops, Canola, Chickpeas, Cotton, Cowpea, Faba beans, Field peas, Lentils, Linola, Linseed, Lupins, Mung beans, Navy beans, Peanuts, Soybeans, Sunflowers and Vetch

DO NOT HARVEST FOR:

Medic and Clover seed crops

7 DAYS AFTER APPLICATION

STOCK FOOD WITHHOLDING PERIODS

DO NOT GRAZE OR CUT FOR STOCK FOR:

Canola, Chickpeas, Cotton, Cowpea, Faba beans, Field peas, Lentils, Linola, Linseed, Lupins, Mung beans, Navy beans, Peanuts, Soybeans, Sunflowers and Vetch

28 DAYS AFTER APPLICATION

Lucerne

21 DAYS AFTER APPLICATION

Medic and Clover pasture

7 DAYS AFTER APPLICATION

COTTON GIN TRASH MUST NOT BE FED TO ANIMALS

GENERAL INSTRUCTIONS

MIXING

- Add water to the spray tank to 10cm above the level of agitation and ensure the agitation device is working vigorously (There must be a minimum of 100 L of water in the tank before any pesticide is added).
- If tank mixing, firstly add any soluble liquid formulations (e.g. Lontrel Herbicide) and allow agitation for approximately 1 minute.
- Then add OzCrop Haloxyfop 520 EC Herbicide at the pot where the agitation is the strongest (do not add OzCrop Haloxyfop 520 EC Herbicide through a strainer or sieve). Allow further agitation for 1 minute.
- Half fill spray tank.
- If using wettable powder or water dispersible granules, or other emulsifiable concentration formulations (e.g. Lorsban 750 WG or Le-Mat), these should be added after the OzCrop Haloxyfop 520 EC Herbicide to the half full spray tank to ensure vigorous agitation.
- Finally add Uptake Spraying Oil or approved alternate spraying oil/wetter (see section on spraying oils/wetters) and continue filling the tank to the required volume maintaining agitation at all times
- Only mix sufficient solution for immediate use. OzCrop Haloxyfop 520 EC Herbicide and any other tank mixes should be applied immediately for best results.

Spraying Oils/Wetters

* **Spraying Oils:** It is essential to add an adjuvant to OzCrop Haloxyfop 520 EC Herbicide. Best results will be achieved with Uptake Spraying Oil at 0.5 L/100 L of spray solution. Alternatively, other oils plus non-ionic wetter may be used. When other crop spraying oils are used, mix at 1.0 L/100 L and add a non-ionic wetter (surfactant) at 200 mL/100 L of spray solution. Use of an oil is not always recommended. See Critical Comments for specific situation recommendations.

† **Non-ionic Wetters:** When Uptake or other oils are not used, a 100% concentrate non-ionic wetting agent such as BS-1000 at 200mL/100L must be used along with the higher rate of OzCrop Haloxyfop 520 EC Herbicide as specified in the Directions for Use.

Where water volumes of less 50 L/ha are used, DO NOT use less than 250 mL/ha of Uptake or 500 mL/ha for oils other than Uptake or less than 100 mL of wetter.

CANOLA, LUCERNE, MEDIC AND CLOVER PASTURES AND SEED CROPS

When tank mixing OzCrop Haloxyfop 520 EC Herbicide with Lontrel herbicides (canola only) or Broadstrike (lucerne, clover and medics), use Uptake Spraying Oil with lower rates of OzCrop Haloxyfop 520 EC Herbicide or a wetting agent with the higher rates of OzCrop Haloxyfop 520 EC Herbicide unless otherwise specified. When mixing OzCrop Haloxyfop 520 EC Herbicide with other broadleaf herbicides on these crops, DO NOT use an oil use a wetter instead.

FIELD PEAS AND CANOLA

The oil recommended is Uptake Spraying Oil. Hasten is also recommended for use with tank-mixtures of OzCrop Haloxyfop 520 EC Herbicide and Select Herbicide.

For canola, OzCrop Haloxyfop 520 EC Herbicide + Lontrel 750SG + Uptake Spraying Oil are compatible and selective to canola. This tank-mixture is also compatible with atrazine or simazine and selective to triazine tolerant canola.

NAVY BEANS, PEANUTS AND SOYBEANS

When mixing with Blazer or Basagran DO NOT add spraying oil to these mixtures. DO NOT use these tank-mixtures on cowpea.

COMPATIBILITY - Ground Use Only

Insecticides: dimethoate, endosulfan, Lorsban 500EC Insecticide, Lorsban 750WG Insecticide, omethoate

Herbicides: atrazine, Basagran, Blazer, Broadstrike Herbicide, Lontrel Herbicide, Lontrel 750SG, MCPA Ester (LVE) – (DO NOT exceed 700 mL/ha of MCPA LVE), oryzalin, Select Herbicide, simazine, Starane 200 Herbicide.

Fungicides: Dithane DF, Dithane Rainshield Fungicide

Trace elements: magnesium sulphate, zinc sulphate

OzCrop Haloxyfop 520 EC Herbicide is not compatible with 2,4-D or MCPA as sodium or amine salts.

Aerial Use

No product other than a recommended crop oil or wetter should be mixed with OzCrop Haloxyfop 520 EC Herbicide when applied by air except for the addition of Lontrel Forestry Herbicide for use in forestry and Lontrel 750SG for use in canola only.

APPLICATION

Apply OzCrop Haloxyfop 520 EC Herbicide in sufficient water to obtain good coverage. It should be applied by an accurately calibrated ground rig or aircraft delivering droplets with a VMD of 200-300 microns.

The following spray volumes are recommended;

Ground Application: 50-150 L/ha

Aerial Application: 30 L/ha minimum

Use higher water volumes in orchards and in dense crops where the weeds may be shielded by the crop canopy.

CLEANING SPRAY EQUIPMENT

Before using OzCrop Haloxyfop 520 EC Herbicide

If broadleaf herbicides, particularly sulfonylureas, have been used in the spray equipment at any time prior to the use of OzCrop Haloxyfop 520 EC Herbicide. Particular care should be taken to follow the directions on the relevant broadleaf herbicide label for equipment cleaning, or damage to susceptible crops may occur.

After using OzCrop Haloxyfop 520 EC Herbicide

Empty the tank and drain the whole system. Thoroughly wash inside the tank using a pressure hose, drain the tank and clean filters in the tank, pump line and nozzles.

To rinse: After cleaning the tank as above, quarter fill the tank with clean water and circulate through the pump, lines hoses and nozzles. Drain and repeat the rinsing procedure twice.

To decontaminate: Before spraying cereals, maize, sorghum or other sensitive crops, wash the tank and rinse the system as above. Then quarter fill the tank and add an alkali detergent (e.g. SURF, Cold Water SURF Concentrate, DynamoMatic Concentrate, OMO or Drive) at 500 mL/100 L of water or the powder equivalent at 500 g/100 L of water, and circulate throughout the system for at least fifteen minutes. Drain the whole system. Remove filters and nozzles and clean separately. Finally flush the system with clean water and allow to drain. Chlorine based cleaners are not recommended.

Rinse water should be discharged onto a designated disposal area, or if this is unavailable, onto unused land away from desirable plants and water sources.

RESISTANT WEEDS WARNING

OzCrop Haloxyfop 520 EC Herbicide is a member of the

GROUP	A	HERBICIDE
-------	----------	-----------

 cyclohexanedione group of herbicides. OzCrop Haloxyfop 520 EC Herbicide has the inhibition of acetyl CoA carboxylase mode of action. For weed resistance management, OzCrop Haloxyfop 520 EC Herbicide is a Group A herbicide. Some naturally occurring weed biotypes resistant to OzCrop Haloxyfop 520 EC Herbicide and other Group A herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by OzCrop Haloxyfop 520 EC Herbicide or other Group A herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to

use, OzCrop Pty Ltd accepts no liability for any losses that may result from the failure of OzCrop Haloxyfop 520 EC Herbicide to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

OzCrop Haloxyfop 520 EC Herbicide damages cereals and grasses. DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible crops/plants, cropping lands or pastures.

Cereal crops or grasses planted within twelve weeks of application may be damaged by the residual effects of OzCrop Haloxyfop 520 EC Herbicide, particularly on light and red soils.

PROTECTION OF LIVESTOCK

DO NOT graze or cut treated crops for stock food except as specified under withholding periods.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

OzCrop Haloxyfop 520 EC Herbicide is toxic to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Do not store near feedstuffs, fertilisers or seeds. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

SMALL SPILL MANAGEMENT

Wear protective equipment (see SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, cat litter or clay granules to the spill. When absorption is complete, sweep up material and contain in a refuse vessel for disposal (see STORAGE AND DISPOSAL section). If necessary wash the spill area with an alkali detergent and water and absorb this wash liquid for disposal as directed above.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening container, preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and washable hat, elbow-length PVC gloves and face shield or goggles. After each day's use, wash gloves and face shield or goggles and contaminated clothing. Wash hands after use.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for OzCrop Haloxyfop 520 EC Herbicide is available from OzCrop on request. Call Customer Service on (02) 8199 3610.

CONDITIONS OF SALE: The use of OzCrop Haloxyfop 520 EC Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by OzCrop Pty Ltd, regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and OzCrop Pty Ltd accepts no responsibility for any consequence whatsoever from the use of this product.

* Other trademarks

Combustible liquid.

In case of fire, use carbon dioxide, dry chemical, foam.

Very toxic to aquatic life with long lasting effects.

Keep away from heat, sparks, open flames and hot surfaces. - No smoking.

Do not eat, drink or smoke when using this product.

IF SWALLOWED: Rinse mouth. Do NOT induce vomiting.



OzCrop Pty Ltd ABN 16 160 656 431

Suite G13/25 Solent Circuit, Norwest, NSW 2153

Tel: 02 8123 0170 • Fax: 02 8123 0171

Email: enquiries@ozcrop.com.au

visit our website www.ozcrop.com.au

APVMA Approval No: 65912/0714

Label Release: 201611

