

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

OzCrop

Metribuzin 750 WG

HERBICIDE

ACTIVE CONSTITUENT: 750 g/kg METRIBUZIN

GROUP **C** HERBICIDE

For selective weed control in cereals, pastures and other crops including vegetables as indicated in the Directions for Use.

NET CONTENTS:

2.5kg, 5kg

NOT A DANGEROUS GOOD
ACCORDING TO THE AUSTRALIAN
DANGEROUS GOODS (ADG) CODE.

BATCH NO.

DATE OF MANUFACTURE:

OzCrop Pty Ltd ABN 16 160 656 431
Suite G13/25 Solent Circuit, Norwest, NSW 2153
Tel: 02 8123 0170 • Fax: 02 8123 0171
Email: enquiries@ozcrop.com.au
www.ozcrop.com.au

DIRECTIONS FOR USE

RESTRAINTS: DO NOT apply to crops that are stressed by any cause (such as severe weather conditions, waterlogging, drought, excessive soil acidity or alkalinity, poor nutrient status, disease, nematode or insect damage) as crop injury may result.

CROP / SITUATION	WEEDS CONTROLLED	STATE	RATE / HA	CRITICAL COMMENTS
Barley	Capeweed, Cat's Ear, Charlock, Chickweed, Common Cotula, Corn Gromwell (Sheepweed, White Ironweed), Corn Spurry, Creeping Speedwell, Crown Beard (Dogweed), Deadnettle (Henbit Lamium spp.) Dock seedlings, Fat Hen, Field Madder, Fumitory, Hare's Ear (Treacle Mustard), Heliotrope, Hogweed (Wireweed), Horehound, Indian Hedge Mustard, Lesser Swinecress, Mouse-ear Chickweed, Mountain Sorrel, (Purple Calandrinia), Rough Poppy, Scarlet Pimpernel, Scrub Nettle (Stinging Nettle), Shepherds Purse, Sorrel seedlings, Sowthistle, Spiny Emex (Doublegee, Three cornered Jack(s)), Storksbill, Toad Rush, Wild Radish, Wild Turnip, Winter Grass, Yellow Burrweed	Vic, Tas only	180 g (Light sandy soils, i.e., sandy loams to loamy sands) 280 g (Medium soils, loams, silt plus clay 40-60%) 380 g (Heavy soils, clay loams, silt plus clay over 60%)	Apply to moist soils while weeds are young and actively growing and barley plants are at the 5-leaf stage. DO NOT use on Barley grown on alkaline light sandy soils with pH over 7. DO NOT use on Barley grown on light sandy soils (Tas).
Barley	Buxbaum's Speedwell, Powell's Amaranth, Stagger weed, Sun Spurge, Great Brome	Tas only		
Barley All varieties excluding Morrell	Capeweed, Common Cotula, Doublegee, Fumitory, Indian Hedge Mustard, Toadrush, Wild Radish, Wild Turnip, Wireweed Brome Grass (<i>Bromus diandrus</i> & <i>B. rigidus</i>) Annual Ryegrass, Barley Grass	WA only Sandy - sandy loam soils	135-150 g plus OzCrop Trifluralin 480 EC at 1.0 L/ha and incorporated by sowing (IBS)	DO NOT apply to Morell barley. Use the lower rate in lighter soils, and the higher rate in heavier soils. Apply to moist soil for best efficacy Seed below herbicide band at 5 cm. Reduced control may occur under high weed burdens (>100/m ²). Ensure OzCrop Metribuzin 750 WG Herbicide is fully dispersed in the boom spray tank prior to the addition of OzCrop Trifluralin 480 EC. Refer to the application section of this and the OzCrop Trifluralin 480 EC label for complete instructions on product use.
Chick peas, Faba Beans, Lentils, Vetch	Refer to the barley section	Vic, SA, WA only	180 g (Light sandy soils, i.e., sandy loams to loamy sands) 280 g (Medium soils, loams, silt plus clay 40-60%) 380 g (Heavy soils, clay loams, silt plus clay over 60%)	Crop: ONLY apply post-sowing, pre-emergence at least 5 cm deep. DO NOT apply post-emergence, as crop injury will occur. Weeds: Weeds should be from pre-emergence to 3 leaf stage except Wireweed (Hogweed) that should not be beyond the cotyledon stage.
Lupins	Capeweed, Doublegee, Wild Radish	WA only	100-150 g plus 100 mL diflufenican	For use on varieties Gunurru, Meritt, Myallie and Kayla only. DO NOT use on Danja, Kiev mutant or Albus lupins. Suppression of larger Doublegee only. Wild Radish weed size up to 250 mm only. Apply to the actively growing lupins from the 3-4 leaf stage onwards. Use higher rates for heavy weed burden and larger weed size. Apply before weeds are covered by crop canopy. DO NOT apply wetting agents or spay oils with this mixture. Some transient leaf burn may occur after application, but this will not affect yield. Best results are obtained if good soil moisture exists at, and is received soon after application. Results may be less effective south of the Great Eastern Highway. Refer to critical comments section on the diflufenican label.

CROP / SITUATION	WEEDS CONTROLLED	STATE	RATE / HA	CRITICAL COMMENTS
Peas (Field and Green) (Not Snow Peas)	Refer to Barley section plus: Buchan Weed, Ball Mustard, Fog Grass (SA only), Dwarf Nettle (Tas only) Suppression only: Sub Clover and Docks (Broadleaved Dock – Tas only), Wild Oats	NSW, Vic, Tas, SA only	180 g (Light sandy soils, i.e., sandy loams to loamy sands)	Crop: Sow crop 3-4cm deep. Apply after rain when crop is pre-emergence to 3-node stage. For post-sowing pre-emergence application allow soil to settle for 2-3 days after sowing before spaying. For post-emergence apply to moist soil within 6 weeks of sowing while weeds are young and actively growing. DO NOT spray crops affected with collar or stem rot. Weeds: Weeds should be from pre-emergence to 3-leaf stage except Wireweed (Hogweed) that should not be beyond the cotyledon stage. Best results for grass and Wireweed control are obtained when rain or irrigation follows within a few days of spraying.
			280g (Medium soils, loams, silt plus clay 40-60%)	
			380 g (Heavy soils, clay loams, silt plus clay over 60%)	
		Qld only	470 g	Apply to medium soils
			700g	Apply to soils high in clay or organic matter content when grass weeds are a problem.
Green Peas (Post emergence) (NOT Snow Peas)		WA only	370-470 g	Refer to Critical Comments for Peas (Field and Green).
Potatoes		ACT, NSW, Vic, Tas, SA, WA only	470 g	Apply after rain or irrigation when crop has no more than 6 leaves or is no more than 10 cm high (many plants may be smaller). CROPS HILLED OR MOLDED UP – Hill or mold up as the crop emerges. Apply 1-4 weeks later as soon as weed seedlings have emerged. If surface crusting is not a problem no further cultivation should be necessary. CROPS NOT HILLED AT OR SOON AFTER PLANTING – Spray when weeds seedling have emerged. No cultivation should be necessary until crop is ready for final hilling. Additional cultivation may be necessary when surface crusting is a problem.
			700 g	Where summer grasses are a problem.
		Vic, Bellarine Peninsula only	280-370 g	Apply to winter crops on light soils only.
		Qld only (Winter crop)	470 g	Apply at first emergence of potatoes to control broadleaf weeds.
		South Qld only (Summer crop)	470-950 g	Where summer grasses such as <i>Echinochloa crus-gali</i> , <i>Eleusine</i> , <i>Digitaria</i> and <i>Panicum</i> spp. are a problem.
		Atherton Tablelands (Qld only)	950 g	Apply to summer crops at first emergence of potatoes to control broadleaved weeds and more troublesome grasses such as <i>Eleusine</i> , <i>Digitaria</i> and <i>Panicum</i> spp.

CROP / SITUATION	WEEDS CONTROLLED	STATE	RATE / HA	CRITICAL COMMENTS
Irrigated Soybeans	Apple of Peru, Cobbler's Pegs, Pigweed, Noogoora Burr, Common Thornapple, Summer Grass, Caltrop	NSW, ACT, NT only	500 g	Apply pre-plant incorporated. Where grass weeds other than summer grass are a problem apply in combination with Ozcrop Trifluralin 480 EC. (Refer to label for directions for use and incorporation techniques). Use only on soils classified as silty clay-heavy clay.
Irrigated Soybeans (Variety Davis only)	Refer to weeds listed for soybeans plus: Anoda Weed, Awnless Barnyard Grass Suppression only of Bellvine	Qld, NT only	750 g	Apply pre-plant incorporated. Where grass weeds other than summer grass are a problem apply in combination with Ozcrop Trifluralin 480 EC. (Refer to label for directions for use and incorporation techniques). Use only on soils classified as silty clay-heavy clay.
Tomatoes Inter-row spaces of furrow irrigated tomatoes	Refer to Peas section plus: Barnyard Grass, Summer Grass, Crowsfoot Grass, Blackberry Nightshade, Pigweed, Black Pigweed, Bull's Head, Green Summer Grass		950 g or 9.5 g/100 m of row	Apply in a 1 metre band centred on the interrow either during or shortly after the formation of the final furrow. Apply to soil moist at the surface. DO NOT exceed spraying pressures of 200 kPa, as excessive drift will occur.
Wheat (Blade only)	Capeweed, Common Cotula, Doublegee, Fumitory, Indian Hedge Mustard, Toad Rush, Wild Turnip, Wild Radish, Wireweed Brome Grass (<i>Bromus diandrus</i> and <i>B. rigidus</i>) Annual Ryegrass, Barley Grass (refer to critical comments)	WA – Great northern region only Sandy - sandy loam soils	200 g and incorporated by sowing (IBS) 200 g/ha plus a OzCrop Trifluralin 480 EC at 1 L/ha and incorporated by sowing (IBS)	OzCrop Metribuzin 750 WG Herbicide has increased crop safety and gives more reliable weed control when incorporated by sowing and trailing harrows. Reduced grass control may occur under high weed burdens (>100/m ²). To achieve good control of Annual Ryegrass and Barley Grass add OzCrop Trifluralin 480 EC at 1 L/ha. Ensure OzCrop Metribuzin 750 WG Herbicide is fully dispersed in the boom spray tank prior to the addition of OzCrop Trifluralin 480 EC. Refer to the application section of this and the OzCrop Trifluralin 480 EC label for complete instructions on product use.
Wheat and Barley Oats	Toad Rush	NSW, ACT, Vic, SA only Vic only	100 g	Toad Rush should be sprayed at the 2-4 leaf stage for optimum control. Spray when soil moisture is plentiful and soil is moist to the surface. Take advantage of dew on the soil surface.
White Lupins	Buchan Weed, Capeweed, Charlock, Chickweed, Deadnettle (<i>Lamium</i> spp.), Fat Hen, Fumitory (Common and Wall Fumitory – Tas), Wireweed (Hogweed), Lesser Swinecress, Purple Calandrinia, Scarlet Pimpernel, Spiny Emex, Corn Spurry (Spurry – Tas), Stinging Nettle (Dwarf Nettle – Tas), Toad Rush, Winter Grass, Heliotrope, Wild Oats, Wild Turnip, Wild Radish Suppression only: Sub Clover and Docks (Broadleaved Dock – Tas)	Tas only	400 g	Spring application – Apply after sowing but before crop emerges.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

TOMATOES: DO NOT APPLY LATER THAN 21 DAYS BEFORE HARVEST.
BARLEY: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION.
OTHER CROPS: NOT REQUIRED IF USED AS DIRECTED.

GENERAL INSTRUCTIONS

OzCrop Metribuzin 750 WG Herbicide is non-corrosive to equipment, non-flammable and non-volatile. The degree of weed control and duration of effect will vary with the amount of herbicide applied, soil texture, rainfall and other conditions.

RESISTANT WEEDS WARNING

OzCrop Metribuzin 750 WG Herbicide is a member of the Triazines group of herbicides. OzCrop Metribuzin 750 WG Herbicide has the inhibitors of photosynthesis at Photosystem II mode of action. For weed resistance management OzCrop Metribuzin 750 WG Herbicide is a Group C Herbicide. Some naturally-occurring weed biotypes resistant to OzCrop Metribuzin 750 WG Herbicide and other Group C Herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides used repeatedly. These resistant weeds will not be controlled by OzCrop Metribuzin 750 WG Herbicide or other Group C herbicides.

GROUP C HERBICIDE

Annual Ryegrass biotypes resistant to diclofop-methyl and other ryegrass specific herbicides may be resistant to OzCrop Metribuzin 750 WG Herbicide. OzCrop Pty Ltd recommends resistance testing of the population to ensure susceptibility prior to use.

Since the occurrence of resistant weeds is difficult to detect prior to use, OzCrop Pty Ltd accepts no liability for any losses that may result from the failure of OzCrop Metribuzin 750 WG Herbicide to control resistant weeds. Large numbers of healthy surviving weeds can be an indication that resistance is developing. Efforts should be made to prevent seed set of the survivors.

DO NOT make more than one application of a Group C Herbicide to a crop in the same season, either pre-sowing incorporated by sowing or post crop and weed emergence. If the user suspects that Group C Herbicide resistant weeds are present, OzCrop Metribuzin 750 WG Herbicide or other Group C Herbicides should not be used.

Strategies to minimise the risk of herbicide resistance are available. Consult your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries.

CROP SAFETY

DO NOT apply to crops that are stressed by any cause (such as severe weather conditions, waterlogging, drought, excessive soil acidity or alkalinity, poor nutrient status, disease, nematode or insect damage) as crop injury may result.

Barley	DO NOT apply to crops grown on sandy soils or soils low in organic matter.
Peas (field & green)	DO NOT apply to Laura, Pennant or White Brunswick varieties as phytotoxicity may occur. DO NOT apply any later than the one node stage to Alma and Dinkum peas. DO NOT spray pea crops in soils or areas where Collar or Stem Rot is likely to occur.
Potatoes	DO NOT apply to crops grown on sandy soils as injury to the crops may result, appearing as chlorosis or minor necrosis. DO NOT apply to Kurell potatoes as phytotoxicity may occur. DO NOT apply to Alantic and Ponitiac varieties – WA only. DO NOT use on dry-grown, non-irrigated crops.
Soybeans	DO NOT use on varieties Cannapolis, Hill and Semstar as crop injury may occur. DO NOT use on soils with a calcareous surface layer as crop injury may occur. DO NOT apply over atrazine residues as crop injury may result. DO NOT apply in combination with soil applied organophosphate insecticides as crop injury may result. DO NOT apply to soils with less than 0.5% organic matter as crop injury may result. DO NOT apply to dryland soybeans. DO NOT apply to soils of pH greater than 8 that are prone to waterlogging as crop injury may occur.
Lupins	DO NOT apply to varieties Danja, Kiev Mutant or Albus Lupins. DO NOT apply to Lupin crops that have received 2.5 litres or more of Simazine pre-seeding, or lupin crops that are showing symptoms of Simazine or Atrazine damage.

MIXING AND APPLICATION COMPATIBILITY

OzCrop Metribuzin 750 WG Herbicide is compatible with the herbicides: chlorsulfuron, metsulfuron, glyphosate, pendimethalin, propargite, diflufenican and OzCrop Trifluralin 480 EC.

MIXING

General

OzCrop Metribuzin 750 WG Herbicide is to be mixed with water and applied as a spray. Before spraying calibrate equipment to determine the quantity of water necessary to uniformly cover the measured area to be treated. Partially fill the tank with water. Measure the amount of herbicide required for the area to be sprayed, adding this amount of OzCrop Metribuzin 750 WG Herbicide to the spray tank with the agitation system engaged. Top up to the correct volume with water.

THE MATERIAL MUST BE KEPT IN SUSPENSION AT ALL TIMES BY CONTINUOUS AGITATION.

In tank mixes OzCrop Metribuzin 750 WG Herbicide must be in suspension before adding the companion herbicide or liquid fertiliser.

Pre-planting Incorporation Treatment:

OzCrop Metribuzin 750 WG Herbicide plus OzCrop Trifluralin 480 EC:

Add OzCrop Metribuzin 750 WG Herbicide to water in spray tank with agitator running as above. Maintain agitation then add OzCrop Trifluralin 480 EC at the recommended rates (Refer to OzCrop Trifluralin 480 EC label).

APPLICATION

General

Best results are obtained if treatment is made to moist soil and moisture is supplied by rainfall or sprinkler irrigation (6-12mm) within 2 weeks after application. If heavy rains occur soon after application, injury to the crop may result.

Aerial Application – (Qld only)

Peas – Apply in a minimum of 20 litres water per hectare.

Potatoes – Apply in a minimum of 50 litres water per hectare. Avoid overlapping of spray swath, Avoid spraying in still conditions and in wind likely to cause drift.

Pre-emergence and Post-emergence (to crop)

OzCrop Metribuzin 750 WG Herbicide should not be applied when weeds are in the young seedling stage (no more than 3 true leaves). Under dry conditions there is no root uptake of OzCrop Metribuzin 750 WG Herbicide and results against grass weeds and Wireweed (Hogweed) will be poor. Except for small areas, use fixed boom power sprayers set at 200-300 kPa (for tomatoes do not exceed 200kPa) and fitted with fan nozzles. Apply a minimum of 80 L-100 L prepared spray/ha. Openings in screens should be equal to or larger than 50 mesh. Agitate by mechanical or hydraulic means in spray tank. When spraying avoid overlapping. Shut off spray booms whilst starting turning, slowing or stopping otherwise crop injury may result.

Pre-plant incorporated – Soybeans only

OzCrop Metribuzin 750 WG Herbicide alone or in combination with OzCrop Trifluralin 480 EC.

Spray anytime from 4 week before sowing until just before sowing. Use fixed boom power sprayers set at 200-300 kPa and fixed with fan nozzles. Use 120-500 litres water per hectare. Openings in screens should be equal to or larger than 80 mesh. Agitate by mechanical or hydraulic means in the spray tank. When spraying avoid overlapping. Shut off spray booms while starting, turning, slowing and stopping otherwise crop injury may result.

OzCrop Metribuzin 750 WG Herbicide alone

If possible spray and incorporate within the same operation. If not possible incorporate spray within 24 hours of application using Go-devil or Lilliston cultivators set to work 5 cm deep.

OzCrop Metribuzin 750 WG Herbicide plus OzCrop Trifluralin 480 EC

After spraying mix soil within 4 hours. Spray and incorporate during the one operation if possible. (Refer to OzCrop Trifluralin 480 EC label for specific incorporation details).

SPRAYER CLEANUP

Thoroughly clean all traces of OzCrop Metribuzin 750 WG Herbicide and /or OzCrop Trifluralin 480 EC from application equipment immediately after use. Flush tank, pump, hoses and boom with several changes of water after removing nozzle tips and screens (clean these parts separately).

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply (except as recommended for crop use) or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots, as injury or loss may occur. DO NOT plant treated area to sensitive crops such as brassicas, sunflowers, beetroot, cucurbits, lettuce or onions for at least 6 months following application of OzCrop Metribuzin 750 WG Herbicide as crop injury may result. Thoroughly work the soil prior to planting of these crops. DO NOT apply under meteorological conditions or from spraying equipment, which could be expected to cause spray to drift onto nearby susceptible plants, adjacent crops, croplands or pastures.

Tomatoes: DO NOT plant follow-on crops of cucurbits or capsicums when applying OzCrop Metribuzin 750 WG Herbicide to autumn plantings.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways by cleaning of equipment, disposal of wastes or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, well-ventilated area, as cool as possible. DO NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulation. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

Harmful if swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. DO NOT inhale dust or spray mist. Wash hands after use.

Safe Work Australia:

Very toxic to aquatic life with long lasting effects.

Do not eat, drink or smoke when using this product.

Avoid release to the environment.

Use personal protective equipment as required.

Brush off loose particles from skin.

Take off contaminated clothing and wash before reuse.

IF SWALLOWED: Rinse mouth. Do NOT induce vomiting.

In case of fire, use carbon dioxide, dry chemical, foam, water fog. Water fog or fine spray is the preferred medium for large fires.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

SDS

Additional information is listed in the safety data sheet (SDS).
A safety data sheet for OzCrop Metribuzin 750 WG Herbicide is available from OzCrop on request. Call Customer Service on (02) 8199 3610.

CONDITIONS OF SALE: The use of OzCrop Metribuzin 750 WG Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by OzCrop Pty Ltd, regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and OzCrop Pty Ltd accepts no responsibility for any consequence whatsoever from the use of this product.



OzCrop Pty Ltd ABN 16 160 656 431

Suite G13/25 Solent Circuit, Norwest, NSW 2153

Tel: 02 8123 0170 • Fax: 02 8123 0171

Email: enquiries@ozcrop.com.au

visit our website www.ozcrop.com.au

APVMA Approval No.: 65831/52134

Label Release: 201707