

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

OzCrop

Propaz 500 WG

HERBICIDE

ACTIVE CONSTITUENT: 500 g/kg PROPYZAMIDE

GROUP **D** HERBICIDE

For selective control of certain grasses and broad leaf weeds in Lettuce, Sports turf, Home lawns and Legume seed crops and pastures, as per the DIRECTIONS FOR USE TABLE.

CONTENTS:

500g, 1kg, 10kg, 15kg, 20kg

NOT A DANGEROUS GOOD
ACCORDING TO THE AUSTRALIAN
DANGEROUS GOODS (ADG) CODE.

BATCH NO.

DATE OF MANUFACTURE:

OzCrop Pty Ltd ABN 16 160 656 431
Suite G13/25 Solent Circuit, Norwest, NSW 2153
Tel: 02 8123 0170 • Fax: 02 8123 0171
Email: enquiries@ozcrop.com.au
www.ozcrop.com.au

DIRECTIONS FOR USE

SITUATION	WEEDS CONTROLLED	STATE	RATE L/HA	CRITICAL COMMENTS	
Lettuce (direct sown)	Barnyard Grass, Blackberry Nightshade, Chickweed, Portulaca, Winter Grass, Rye Grass, Summer Grass,	All States	4.5 kg in 330-550 L	Apply pre-emerged, immediately after sowing. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours.	
Transplanted	Wireweed, Prince of Wales Feather, Shepherd's Purse, Nettles			Apply immediately after transplanting. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours.	
		Vic only	2.3 kg in 300-550 L	Apply on light sandy soils, low in organic matter.	
Turf & Lawns	Winter Grass (<i>Poa annua</i>)	All States	1.2 kg (12g/100m ²)	RATE/ BOWLING GREEN 37m x 37m	Apply during the Autumn and Winter, if established turf is infested with seedling winter grass. Ensure a uniform coverage. Irrigate immediately after application with 12 – 25 mm irrigation or rain and again 24 hour later. Repeat treatment if necessary on Qld blue couch after 4 weeks. DO NOT RESEED FOR 60 DAYS AFTER TREATMENT. Use ONLY on established turf. DO NOT use on Bluegrass, Ryegrass, Fescue or Bent grass lawns.
Common Couch				170g	
Queensland Blue Couch		Qld only	0.6 kg (6 g/100m ²)	85g	
Buffalo Grass		WA only	1.2 kg (12g/100m ²)	170g	
Lucerne, Clover, Medics etc. Grown for Forage, Hay or Seed production	Annual Ryegrass, Winter Grass, Barley Grass, Silver Grass, Prairie Grass, Canary Grass, Great Brome, Squirrel Tail Fescue, Rats Tail Fescue	NSW, ACT, Tas, SA, WA only	1-1.5 kg	SEEDLING CROPS: Apply after the third trifoliolate leaf is fully developed. ESTABLISHED CROPS: Apply any time before the end of July. WEEDS: Apply when weeds are at post- or pre-emergent stages. BEST RESULTS: When applied during cool and moist conditions. Rain or irrigation is necessary to thoroughly wet the top 3 to 4 cm of soil. This is similar to 25mm rain. Use rates towards the higher end of the range on heavy soils, if conditions are not optimal or where a heavy grass population is expected. Grazing or cutting PRIOR to treatment is recommended.	
	Wild oats, Perennial Ryegrass, Bent grass, English couch, Yorkshire fog grass, Chickweed, Seedling Sorrel, Blackberry nightshade, Patterson's curse, seedling Wireweed.		1.5-2 kg		

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS: DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD OR FOR SEED FOR 25 DAYS AFTER APPLICATION.

NOT CLASSIFIED AS DANGEROUS GOODS FOR TRANSPORT BY ROAD AND RAIL

GENERAL INSTRUCTIONS

Best results are obtained by applying the product to cool, wet soil.

Spray Preparation

OzCrop Propaz 500 WG Herbicide is a water dispersible granule formulation and must be thoroughly dispersed in water and applied as a dilute spray solution.

- Partly fill the boomspray-tank with water so to form a slurry.
- Add the required amount of OzCrop Propaz 500 WG Herbicide for the area to be sprayed into water containers.
- Add this correct amount of pre-mixed OzCrop Propaz 500 WG Herbicide into the boomspray-tank with the agitation system operating.

- Fill up to the correct volume with water. The mixture must be kept in suspension at all times by continuous agitation, otherwise bottom settling of the product will result. Re-dispersion of the product will be extremely difficult and blockages of filters are likely to occur.
- When mixing with other chemical products ensure to check all label instructions prior to chemicals being tank mixed with OzCrop Propaz 500 WG Herbicide to check for chemical compatibility. Test a small area for any adverse affects prior to application.

DO NOT mix with any products that contains the active constituent, MCPA. If tank mixing, OzCrop Propaz 500 WG Herbicide must be in suspension before adding other herbicides, insecticides or surfactant. Check all labels Directions for Use table prior to tank mixing with OzCrop Propaz 500 WG Herbicide . Wash sprayer thoroughly after use.

Equipment

Always ensure the sprayer is clean, using a suitable spray and boom cleaner as directed. It is recommended that for turf situations, up to 100 L/ha spray volume is required.

RESISTANT WEEDS WARNING

OzCrop Propaz 500 WG Herbicide is a member of the benzamide group of herbicides with the inhibitors of microtubal assembly mode of action group of herbicides. For weed resistance management this is a Group D herbicide. Some naturally occurring weed biotypes resistant to OzCrop Propaz 500 WG Herbicide and other amide herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by OzCrop Propaz 500 WG Herbicide or other amide herbicides.

GROUP D HERBICIDE

Since the occurrence of resistant weeds is difficult to detect prior to use, OzCrop Pty Ltd accepts no liability for any losses that may result from the failure of OzCrop Propaz 500 WG Herbicide to control the resistant weeds. Advice as to strategies and alternative treatments that can be used should be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or a OzCrop representative.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

PROTECTION OF LIVESTOCK

DO NOT graze sheep, cut for stock feed (including poultry and pigs) any treated areas (for 25 days after application).

DO NOT FEED TURF/LAWN CLIPPINGS TO POULTRY OR LIVESTOCK.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply to, or allow spray to drift onto crops, pastures or non target – desirable plants, nor onto agricultural land scheduled for cropping or pasture, as injury to the crop or pasture may occur.

STORAGE AND DISPOSAL

- Keep out of reach of children.
- Store in the closed, original container in a dry, cool, well-ventilated area, out of direct sunlight. Keep from contact with fertilisers, insecticides, fungicides and seeds.
- DO NOT re-use container or dispose of undiluted chemicals on-site.
- DO NOT dispose of undiluted chemicals on-site. Puncture or shred and bury empty containers in a local authority landfill. If not available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation, and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. Avoid inhaling the dust. When preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, washable hat and elbow length PVC gloves. After use, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

Safe Work Australia:
May cause respiratory irritation.
Suspected of causing cancer.
Toxic to aquatic life with long lasting effects.
IF SWALLOWED: Rinse mouth. Do NOT induce vomiting.
IF ON SKIN: Wash with plenty of soap and water.
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
If skin / eye irritation occurs and persists: Get medical advice.
In case of fire, use carbon dioxide, dry chemical, foam, water fog.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

SDS

Additional information is listed in the safety data sheet (SDS). A safety data sheet for OzCrop Propaz 500 WG Herbicide is available from OzCrop on request. Call Customer Service on (02) 8199 3610.

CONDITIONS OF SALE: The use of OzCrop Propaz 500 WG Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by OzCrop Pty Ltd, regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and OzCrop Pty Ltd accepts no responsibility for any consequence whatsoever from the use of this product.



OzCrop Pty Ltd ABN 16 160 656 431

Suite G13/25 Solent Circuit, Norwest, NSW 2153

Tel: 02 8123 0170 • Fax: 02 8123 0171

Email: enquiries@ozcrop.com.au

visit our website www.ozcrop.com.au

APVMA Approval No.: 68438/58369

Label Release: 201710

