

# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

 OzCrop

# Propaz

**HERBICIDE**

ACTIVE CONSTITUENT: 500 g/L PROPYZAMIDE  
SOLVENT: 42 g/L ETHYLENE GLYCOL

GROUP **D** HERBICIDE

*For selective control of certain grasses and broad leaf weeds in Canola, Chicory, Endive, Lettuce, Radicchio, Sports turf, Home Lawns and Legume seed crops and pastures as per the Directions for Use Table. Also for the control of annual ryegrass in Pyrethrum crops.*

CONTENTS:

**5L -1000L**

NOT A DANGEROUS GOOD  
ACCORDING TO THE AUSTRALIAN  
DANGEROUS GOODS (ADG) CODE.

BATCH NO.

DATE OF MANUFACTURE:

**OzCrop Pty Ltd** ABN 16 160 656 431  
G13/25 Solent Circuit, NORWEST NSW 2153  
Tel: 02 8123 0170 • Fax: 02 8123 0171  
Email: [enquiries@ozcrop.com.au](mailto:enquiries@ozcrop.com.au)  
[www.ozcrop.com.au](http://www.ozcrop.com.au)

**DIRECTIONS FOR USE**

SITUATION	WEEDS CONTROLLED	STATE	RATE L/HA	CRITICAL COMMENTS	
Canola and Mustard (oilseed cultivars, Brassica juncea)	Annual ryegrass, Barley grass, Canary grass, Fescue, Great Brome, Prairie grass, Rat's tail fescue, Silver grass, Squirrel tail fescue, Wild oats, Winter grass	All States	1	Incorporate by sowing (IBS) when weeds are at the pre-emergent stage.	
Lettuce (direct sown)	Barnyard Grass, Blackberry Nightshade, Chickweed, Portulaca, Winter Grass, Rye Grass, Summer Grass, Wireweed, Prince of Wales Feather, Shepherd's Purse, Nettles	All States	4.5 in 330-550 L	Apply pre-emerged, immediately after sowing. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours.	
Transplanted				Apply immediately after transplanting. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours.	
			Vic only	2.3 in 300-550 L	Apply on light sandy soils, low in organic matter.
Turf & Lawns	Winter Grass ( <i>Poa annua</i> )  Common Couch  Queensland Blue Couch  Buffalo Grass	All States	1.2 (12 mL/100m <sup>2</sup> )	<b>RATE BOWLING GREENS 37m x 37m</b> Apply during the Autumn and Winter, if established turf is infested with seedling Winter Grass.	
				170 mL	Ensure a uniform coverage. Irrigate immediately after application with 12-25 mm irrigation or rain and again 24 hours later.
		Qld only	0.6 (6 mL/100m <sup>2</sup> )	85 mL	
		WA only	1.2 (12 mL/100m <sup>2</sup> )	170 mL	Repeat treatment if necessary on Queensland Blue Couch after 4 weeks. <b>DO NOT RESEED FOR 60 DAYS AFTER TREATMENT.</b> Use only on established turf. DO NOT use on Bluegrass, Ryegrass, Fescue or Bent grass lawns.
Lucerne, Clover, Medics etc.  Grown for Forage, Hay or Seed production	Annual Ryegrass, Winter Grass, Barley Grass, Silver Grass, Prairie Grass, Canary Grass, Great Brome, Squirrel Tail Fescue, Rats Tail Fescue	NSW, ACT, Tas, SA, WA only	1-1.5	SEEDLING CROPS: Apply after the third trifoliolate leaf is fully developed. ESTABLISHED CROPS: Apply any time before the end of July. WEEDS: Apply when weeds are at post- or pre-emergent stages. BEST RESULTS: When applied during cool and moist conditions. Rain or irrigation is necessary to thoroughly wet the top 3 to 4 cm of soil. This is similar to 25mm rain. Use rates towards the higher end of the range on heavy soils, if conditions are not optimal or where a heavy grass population is expected. Grazing or cutting prior to treatment is recommended.	
	Wild Oats, Perennial Ryegrass, Bent Grass, English Couch, Yorkshire Fog Grass, Chickweed, seedling Sorrel, Blackberry Nightshade, Patterson's Curse, seedling Wireweed		1.5-2		
Pyrethrum crops ( <i>Tanacetum cinerariifolium</i> )	Annual ryegrass	All States	2.5-4.5 apply in a spray volume of 330-550L/ha	Apply only in crops where there is evidence of Group A or B herbicide resistant ryegrass (HRR). Apply a maximum of 2 foliar applications per season at a minimum interval of 30 days. Crop may be treated either at pre-emergence stage, or at post-emergence stage any time after the third leaf has fully developed.  Best results are obtained when the product is applied during cool moist conditions. Rain or irrigation is necessary to thoroughly wet the top 3-4cm of soil. This is similar to 23 mm of rain. Use low rate for light, sandy type soils. Use higher rate on heavier soils, if conditions are not optimal or where a heavy grass population is expected. Grazing or cutting of crop before treatment is recommended.	

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD**

**HARVEST:**

**Canola, Chicory, Endive, Mustard, Radicchio: NOT REQUIRED WHEN USED AS DIRECTED**

**GRAZING:**

**Chicory, Endive, Radicchio: DO NOT GRAZE OR CUT FOR STOCK FOOD**

**Pyrethrum crops: DO NOT GRAZE OR CUT TREATED AREA FOR STOCK FOOD FOR 25 DAYS AFTER APPLICATION.**

**Canola, Mustard: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 12 WEEKS AFTER APPLICATION.**

**All other crops: DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD OR FOR SEED FOR 25 DAYS AFTER APPLICATION.**

## GENERAL INSTRUCTIONS

Best results are obtained by applying the product to cool, wet soil.

## MIXING

Prior to pouring, shake container vigorously, then add the required quantity of OzCrop Propaz Herbicide to water in the spray vat while stirring or with agitators in motion. DO NOT mix with other pesticides.

## RESISTANT WEEDS WARNING

OzCrop Propaz Herbicide is a member of the benzamide group of herbicides. The product has the inhibitors of microtubule assembly mode of action. For weed resistance management OzCrop Propaz Herbicide is a Group D herbicide. Some naturally-occurring weed biotypes resistant to OzCrop Propaz Herbicide and other Group D herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by OzCrop Propaz Herbicide or other Group D herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, OzCrop Pty Ltd accepts no liability for any losses that may result from failure of OzCrop Propaz Herbicide to control resistant weeds. Advice as to strategies and alternative treatments that can be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or OzCrop representative.

## PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

## PROTECTION OF LIVESTOCK

DO NOT graze sheep on treated areas or feed to poultry or pigs.

## PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, which may cause spray drift onto nearby susceptible plants, crops, cropping land or pastures.

## STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area, out of direct sunlight. Keep away from contact with fertilisers, insecticides, fungicides and seeds. Protect from frost. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

**For refillable containers**, empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

**GROUP D HERBICIDE**

## SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. When preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, washable hat and elbow-length PVC gloves. After use, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

### Safe Work Australia:

Suspected of causing cancer.

Very toxic to aquatic life with long lasting effects.

Avoid release to the environment.

IF SWALLOWED: Rinse mouth. Do NOT induce vomiting.

IF ON SKIN: Wash with plenty of soap and water.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

If exposed or concerned: Get medical advice.

If skin irritation occurs: Get medical advice.

If eye irritation persists: Get medical advice.

Not combustible. Use extinguishing media suited to burning materials.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126, New Zealand 0800 764 766.

## SAFETY DATA SHEET

Additional information is listed in the safety data sheet (SDS). A safety data sheet for OzCrop Propaz Herbicide is available from OzCrop on request. Call Customer Service on (02) 8123 0170.

**CONDITIONS OF SALE:** The use of OzCrop Propaz Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by OzCrop Pty Ltd, regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and OzCrop Pty Ltd accepts no responsibility for any consequence whatsoever from the use of this product.



**OzCrop Pty Ltd** ABN 16 160 656 431

G13/25 Solent Circuit, NORWEST NSW 2153

Telephone: (02) 8123 0170 • Facsimile: (02) 8123 0171

Email: [enquiries@ozcrop.com.au](mailto:enquiries@ozcrop.com.au)

visit our website [www.ozcrop.com.au](http://www.ozcrop.com.au)

APVMA Approval No.: 66883/134376

Label Release: 202205

