

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

OzCrop

Quizalofop-P-Ethyl 200 EC

SELECTIVE HERBICIDE

ACTIVE CONSTITUENT: 200 g/L QUIZALOFOP-P-ETHYL
SOLVENTS: 592 g/L HYDROCARBON LIQUID
150 g/L N-METHYL-PYRROLIDONE

GROUP A HERBICIDE

For the control of certain grass weeds in broadleaf crops as per Directions for Use table.

CONTENTS:

10L, 20L, 110L, 200L, 1000L



UN No. 3082
ENVIRONMENTALLY HAZARDOUS
SUBSTANCE, LIQUID, N.O.S.
PACKING GROUP III
HAZCHEM CODE 2X
BATCH NO.
DATE OF MANUFACTURE:

OzCrop Pty Ltd

ABN 16 160 656 431
Suite G13/25 Solent Circuit, Norwest,
NSW 2153
Tel: 02 8123 0170 • Fax: 02 8123 0171
Email: enquiries@ozcrop.com.au
www.ozcrop.com.au

DIRECTIONS FOR USE

RESTRAINTS:

DO NOT apply to crops that are stressed by any cause (such as severe cold weather, disease, drought or waterlogging).
 DO NOT apply to weeds that are stressed by any cause (such as weather conditions, drought, waterlogging) or not actively growing. Grass weeds under stress frequently become less susceptible to herbicide activity. This may result in an incomplete kill or suppression of weeds only.
 DO NOT apply under conditions of high temperature and low humidity.
 DO NOT apply if rainfall is expected within 3 hours.
 DO NOT apply less than 50 L prepared spray/ha – see section "Ground Spraying".

BOOM SPRAYING: WINTER GROWING WEEDS: ALL STATES

CROPS	WEEDS CONTROLLED	WEEDS STAGE	RATE/ha	WITHHOLDING PERIOD		CRITICAL COMMENTS
				HARVEST	GRAZING	
Beetroot Cabbage Canola Carrots Cauliflower Chickpeas	Annual Ryegrass (<i>Lolium rigidum</i>), Brome Grasses (<i>Bromus</i> spp.)	3 leaf to early tillering	150 mL or 190 mL	14 days 9 weeks 11 weeks 10 weeks 14 days 12 weeks	– – 4 weeks – – 4 weeks	Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*.
Clover & subclover pastures Cucumbers • Faba beans Field peas	Barley Grass (<i>Hordeum</i> spp.), Volunteer Barley, Volunteer Wheat	3 leaf to early tillering	125 mL	– 14 days 12 weeks 9 weeks	14 days – 4 weeks 4 weeks	Apply when weeds are actively growing*.
Honey dew melons • Lentils Lucerne Lupins Medic pasture Mung beans Navy beans Onions Potatoes Pumpkins • Radish Tomatoes Vetch	Wild Oats (<i>Avena</i> spp.)	3 leaf to mid tillering	65 mL or 125 mL	9 weeks 12 weeks – 6 weeks – 12 weeks 12 weeks 18 weeks 10 weeks 9 weeks 21 days 4 weeks 12 weeks	– 4 weeks 4 weeks 4 weeks 14 days 4 weeks 4 weeks – – – – – – 4 weeks	Apply when weeds are actively growing. Use the lower rate only under light weed pressure and/or when weeds are at the 1-3 leaf stage. Use higher rate under heavy weed pressure and/or when weeds are tillering. The higher rate will give better or more reliable control*.

* Always add a surfactant/wetting agent at the recommended rate. See section on "use of surfactant/wetting agent" for details.

• **IMPORTANT:** DO NOT apply OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide before the five true leaf stage of Cucumbers, Honey Dew Melons and Pumpkins.

BOOM SPRAYING: SUMMER GROWING WEEDS: ALL STATES

CROPS	WEEDS CONTROLLED	WEEDS STAGE	RATE/ha	WITHHOLDING PERIOD		CRITICAL COMMENTS
				HARVEST	GRAZING	
Beetroot Cabbage Cauliflower Carrots Cucumbers •	Awnless Barnyard Grass (<i>Echinochloa colona</i>), Barnyard Grass (<i>Echinochloa crusgalli</i>)	3 leaf to early tillering	250 mL or 375 mL	14 days 9 weeks 14 weeks 10 weeks 14 days	– – – – –	Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*.
Green beans Honey dew melons • Lucerne Mung beans Navy beans	Couch Grass (<i>Cynodon dactylon</i>)	2-4 leaf seedling	250 mL	5 weeks 9 weeks – 12 weeks 12 weeks	4 weeks – 4 weeks 4 weeks 4 weeks	Apply when weeds are actively growing. This rate will only control seedling Couch Grass which is less than 2 months old and established from seed*.
Onions Peanuts Potatoes Pineapples Pumpkins • Radish Soybeans Sunflower Tomatoes	Dinebra (<i>Dinebra retroflexa</i>)	3 leaf to early tillering	125 mL or 190 mL	18 weeks 11 weeks 10 weeks 7 days 9 weeks 21 days 12 weeks 9 weeks 4 weeks	– 11 weeks – – – – 4 weeks 11 weeks –	Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*.
	Crowsfoot Grass (<i>Eleusine indica</i>)	2 leaf to early tillering	250 mL			Apply when weeds are actively growing*.



BOOM SPRAYING: SUMMER GROWING WEEDS: ALL STATES – continued

CROPS	WEEDS CONTROLLED	WEEDS STAGE	RATE/ha	WITHHOLDING PERIOD		CRITICAL COMMENTS
				HARVEST	GRAZING	
Beetroot Cabbage Cauliflower Carrots Cucumbers • Green beans Honey dew melons • Lucerne Mung beans Navy beans Onions Peanuts Potatoes Pineapples Pumpkins • Radish Soybeans Sunflower Tomatoes	Foxtail Millet (<i>Setaria italica</i>) Columbus Grass (<i>Sorghum x alnum</i>), Johnson Grass (<i>Sorghum halepense</i>) Liverseed Grass (<i>Urochloa panicoides</i>) Rhodes Grass (<i>Chloris gayana</i>), Stink Grass (<i>Eragrostis cilianensis</i>) Queensland Blue Grass (<i>Dichanthium sericeum</i>) Summer Grass (<i>Digitaria ciliaris</i>)	2 leaf to early tillering 2 leaf to 3 tiller 3 leaf to early tillering 3 leaf to early tillering 3 leaf to early tillering	125 mL or 250 mL 250 mL or 375 mL 125 mL or 250 mL 250 mL 250 mL or 500 mL	14 days 9 weeks 14 weeks 10 weeks 14 days 5 weeks 9 weeks – 12 weeks 12 weeks 18 weeks 11 weeks 10 weeks 7 days 9 weeks 21 days 12 weeks 9 weeks 4 weeks	– – – – – 4 weeks – 4 weeks 4 weeks – – 11 weeks – – – – 4 weeks 11 weeks –	Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*. Apply when weeds are actively growing*. Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering*.

* Always add a surfactant/wetting agent at the recommended rate. See section on “use of surfactant/wetting agent” for details.

- **IMPORTANT:** DO NOT apply OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide before the five true leaf stage of Cucumbers, Honey Dew Melons and Pumpkins.

HAND GUN SPRAYING: ALL STATES

CROPS	WEEDS CONTROLLED	WEEDS STAGE	RATE/ha	WITHHOLDING PERIOD		CRITICAL COMMENTS
				HARVEST	GRAZING	
Beetroot Cabbage Cauliflower Canola Carrots Chickpeas Clover & subclover pastures Cucumbers • Faba beans Field peas Grapes Green beans Honey dew melons • Lentils Lucerne Lupins Medic pasture Mung beans Navy beans Non crop Onions Peanuts Pineapples Potatoes Pumpkins • Radish Soy beans Sunflowers Tomatoes Vetch	Awnless Barnyard Grass (<i>Echinochloa colona</i>), Crowfoot Grass (<i>Eleusine indica</i>) Paspalum (<i>Paspalum dilatatum</i>) Johnson Grass (<i>Sorghum halepense</i>) Kikuyu (<i>Pennisetum clandestinum</i>)	3 leaf to early tillering Seedling to Established Clumps Seedling 2 leaf to early tillering Rhizome 2 leaf to early tillering Seedling to Established plants	65 mL 125 mL	14 days 9 weeks 14 days 11 weeks 10 weeks 12 weeks – 14 days 12 weeks 9 weeks – 5 weeks 9 weeks 12 weeks – 6 weeks – 12 weeks 12 weeks – 18 weeks 11 weeks 7 days 10 weeks 9 weeks 21 days 12 weeks 9 weeks 4 weeks – 12 weeks	– – – 4 weeks – 4 weeks 14 days – 4 weeks 4 weeks – 4 weeks 4 weeks – 4 weeks 14 days 4 weeks 4 weeks – – – 11 weeks – – – 4 weeks 11 weeks – 4 weeks	Apply when weeds are actively growing. Use a minimum of 800L of prepared spray/ha. Thoroughly wet target weeds (especially well established clumps) without causing runoff. Repeat spraying may be necessary for perennial grass species. Always add a surfactant/wetting agent at the recommended rate – see section on “Use of Surfactant/Wetting agent” for details.

- **IMPORTANT:** DO NOT apply OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide before the five true leaf stage of Cucumbers, Honey Dew Melons and Pumpkins.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.



WITHHOLDING PERIODS:

HARVEST

BEETROOT, CAULIFLOWER, CUCUMBERS: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.

CABBAGE, FIELD PEAS, HONEY DEW MELON, PUMPKINS, SUNFLOWERS: DO NOT HARVEST FOR 9 WEEKS AFTER APPLICATION.

CARROTS, POTATOES: DO NOT HARVEST FOR 10 WEEKS AFTER APPLICATION.

CHICKPEAS, FABA BEANS, LENTILS, MUNG BEANS, NAVY BEANS, SOYBEANS, VETCH: DO NOT HARVEST FOR 12 WEEKS AFTER APPLICATION.

CANOLA, PEANUTS: DO NOT HARVEST FOR 11 WEEKS AFTER APPLICATION.

GREEN BEANS: DO NOT HARVEST FOR 5 WEEKS AFTER APPLICATION.

GRAPES: NOT REQUIRED WHEN USED AS DIRECTED.

LUPINS: DO NOT HARVEST FOR 6 WEEKS AFTER APPLICATION.

ONIONS: DO NOT HARVEST FOR 18 WEEKS AFTER APPLICATION.

PINEAPPLES: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.

RADISH: DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION.

TOMATOES: DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION.

GRAZING

CANOLA, CHICKPEAS, FABA BEANS, FIELD PEAS, GREEN BEANS, LENTILS, LUCERNE, LUPINS, MUNG BEANS, NAVY BEANS, SOYBEANS, VETCH: DO NOT GRAZE OR CUT FOR STOCK FEED FOR 4 WEEKS AFTER APPLICATION.

CLOVER, MEDIC PASTURES (EXCEPT LUCERNE): DO NOT GRAZE OR CUT FOR STOCK FEED FOR 14 DAYS AFTER APPLICATION.

PEANUTS, SUNFLOWERS: DO NOT GRAZE OR CUT FOR STOCK FEED FOR 11 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide is an emulsifiable concentrate herbicide used post emergence for the control of certain grass weeds in specified broadleaf crops.

OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide is a systemic herbicide, which is rapidly absorbed and readily translocates from the treated foliage to the root system and growing points of the plant. Applications made to grass weeds stressed due to insufficient moisture or cold temperatures may result in unsatisfactory control.

TIME OF APPLICATION

For maximum benefit, weeds should be sprayed when young and actively growing. Application may be made at any stage of crop growth (except to curcubit crops), but sufficient time must be allowed for the prescribed withholding period to expire before harvest and / or grazing or cutting for stockfeed.

IMPORTANT: DO NOT apply OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide to curcubit crops before the five true leaf stage.

SPRAY PREPARATION

If spray equipment has been previously used for the application of sulfonylurea herbicides (eg. metsulfuron methyl), chlorsulfuron, thifensulfuron methyl, triasulfuron, ensure that the recommended sprayer equipment procedure for the product previously applied has been followed, before using the equipment for the application of OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide. OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide is an emulsifiable concentrate to be mixed with water and applied as a spray. Add the required amount of OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide directly to the spray tank and mix well. If mixing with mancozeb or carbendazim, add these to the spray tank and mix thoroughly PRIOR to adding OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide. If spray solution has been left standing for more than 3 hours ensure thorough re-agitation before commencement of spraying. DO NOT allow spray mixture to stand overnight, as re-suspension may be difficult.

USE OF A SURFACTANT/WETTING AGENT

OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide may be used with either of the following adjuvant system:

- 1) A methylated and/or ethylated seed crop oil (eg Hasten[®]) at 1 L/100 L of final spray volume (i.e. 1% v/v).
- 2) A non-ionic surfactant (1000 gai/L) at 200 mL/100L of final spray volume (i.e. 0.2% v/v).

DO NOT use OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide with mineral based crop oils (eg D-C-Trate[®]) or non-methylated spray oils of crop origin (eg Codacide[®] oil, Synetrol[®] oil) as spray adjuvants.

COMPATIBILITIES

OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide is compatible with the herbicides sethoxydim and clethodim. The recommended surfactant/wetting agent is the methylated and/or ethylated seed crop oil adjuvant system. OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide is compatible with the insecticides deltamethrin, omethoate, dimethoate, methomyl, endosulfan, the fungicides carbendazim and mancozeb and the herbicide clopyralid when using the methylated and/or ethylated seed crop oil adjuvant system. OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide is compatible with the herbicide diflufenican when using the non-ionic surfactant only adjuvant system.

DO NOT add crop oil to tank mixtures of OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide /diflufenican.

Application of OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide /diflufenican tank mix may result in crop effects, which usually appear as yellow or white bleaching on the leaves. New growth and subsequent yield, however, will not be affected.

To minimise crop effects apply OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide a minimum of 5 days prior to diflufenican application or a minimum 10 days after diflufenican application (14-21 days if weeds have been stressed).

GROUND SPRAYING

Use between 50 and 100 litres of prepared spray/ha, the higher spray volume gives more reliable results. If weeds are large and the crop canopy is dense, increase the volume to 150 litres prepared spray/ha.

Use sprayers with flat fan or hollow cone nozzles at 210-350 kPa to give an even distribution of droplets with no less than 50% of the spray volume in the 150-300 micron range. 1100 flat fan nozzles are preferred. For proper spray coverage adjust the boom and nozzle height according to the specification listed by the nozzle manufacturers. Ensure the spray is evenly distributed and a double overlap spray pattern is obtained. Avoid overlapping boomspray widths and shut off spray booms while starting, slowing or stopping.

SPRAYER CLEANUP

Thoroughly clean all traces of OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide from application equipment immediately after use. Flush tank, pump, hoses and boom with several changes of water after removing nozzle tips and screens (clean those parts separately). Failure to thoroughly clean the equipment may result in injury to subsequently sprayed cereal and grass crops.

If spray equipment has been previously used for the application of sulfonylurea herbicides eg (Metsulfuron methyl), thifensulfuron methyl, chlorsulfuron, triasulfuron ensure that the recommended sprayer cleanup procedure of the product previously sprayed has been followed before using the equipment for the application of OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide.

RESISTANT WEEDS WARNING

OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide is a member of the "Fop" (aryloxyphenoxy propionate) group of herbicides. OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide has the inhibitors of acetyl coA carboxylase mode of action. For weed resistance management OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide is a Group A herbicide. Some naturally occurring weed biotypes resistant to OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide and other Group A herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide or other Group A herbicides.

GROUP A HERBICIDE

Since the occurrence of resistant weeds is difficult to detect prior to use, OzCrop Pty Ltd accepts no liability for any losses that may result from failure of OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide to control resistant weeds. Strategies to minimise the risk of herbicide resistance are available. Consult your farm chemical supplier, consultant, OzCrop Representative, local Department of Agriculture or Primary Industries.

PRECAUTION

Re-Entry Period: DO NOT allow entry into treated areas until the spray has dried. When prior entry is necessary, wear chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT plant cereal crops into the treated area for a period of 18 weeks after application of OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide.

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping land or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

10L, 20L and 200L only: DO NOT store tank-mixes of OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide with other materials. Store in the closed original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. When storing, keep from contact with fertilisers, fungicides and seeds. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the container below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

If the container is a returnable/refillable type, when empty, or contents no longer required, return the container to the point of purchase.

Refillable Containers (1000L only): Store in closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

Envirodrum 110L Mini Bulk Returnable Container: Store the original sealed container in a cool well ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with Micro Matic valve or security seal. DO NOT contaminate the container with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the container have been used, please return the empty container to the point of purchase.

SAFETY DIRECTIONS: Will irritate the eyes and skin. Avoid contact with eyes. When opening the container, preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, and elbow-length PVC gloves. If product on skin immediately wash area with soap and water. Wash hands after use. After each day's use wash gloves and contaminated clothing.

Safe Work Australia:
Combustible liquid.
May be fatal if swallowed and enters airways.
Harmful if inhaled.
May cause respiratory irritation.
May damage fertility or the unborn child.
Very toxic to aquatic life with long lasting effects.
Avoid breathing fumes, mists, vapours or spray.
Use only outdoors or in a well ventilated area.
IF SWALLOWED: Immediately call a POISON CENTER or doctor.
IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing.
If skin irritation occurs: Get medical advice.
In case of fire, note the following. Alcohol resistant foam is the preferred firefighting medium but, if it is not available, normal foam can be used. Try to contain spills, minimise spillage entering drains or water courses.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed do NOT induce vomiting. Give a glass of water. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

SDS: Additional information is listed in the safety data sheet (SDS). A safety data sheet for OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide is available from OzCrop on request. Call Customer Service on (02) 8199 3610.

CONDITIONS OF SALE: The use of OzCrop Quizalofop-P-Ethyl 200 EC Selective Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by OzCrop Pty Ltd, regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and OzCrop Pty Ltd accepts no responsibility for any consequence whatsoever from the use of this product.

Not a trademark of OzCrop Pty Ltd

OzCrop Pty Ltd ABN 16 160 656 431

Suite G13/25 Solent Circuit, Norwest, NSW 2153

Tel: 02 8123 0170 • Fax: 02 8123 0171 • Email: enquiries@ozcrop.com.au

visit our website www.ozcrop.com.au

APVMA Approval No.: 68433/58363

Label Release: 201710